

BEST AVAILABLE COPY

PCT/AU2004/001300



REC'D 12 OCT 2004

WIPO

PCT

**PRIORITY
DOCUMENT**

SUBMITTED OR TRANSMITTED IN
COMPLIANCE WITH RULE 17.1(a) OR (b)

**Patent Office
Canberra**

I, JULIE BILLINGSLEY, TEAM LEADER EXAMINATION SUPPORT AND SALES hereby certify that annexed is a true copy of the Provisional specification in connection with Application No. 2003905222 for a patent by QUEENSLAND UNIVERSITY OF TECHNOLOGY as filed on 25 September 2003.



WITNESS my hand this
Sixth day of October 2004

A handwritten signature in cursive script, reading "J. Billingsley".

JULIE BILLINGSLEY
TEAM LEADER EXAMINATION
SUPPORT AND SALES

Regulation 3.2

A U S T R A L I A

Patents Act 1990

PROVISIONAL SPECIFICATION

for the invention entitled:

“Resistance elements and uses therefor”

The invention is described in the following statement:

RESISTANCE ELEMENTS AND USES THEREFOR

FIELD OF THE INVENTION

[0001] THIS INVENTION relates generally to pathogenic resistance. More particularly, the present invention relates to polynucleotide and polypeptide sequences
5 involved in the resistance mechanism of plants to pathogens, especially fungal pathogens. The present invention also relates to the use of these sequences for modulating plant resistance and for producing genetically modified plants having modified pathogen resistance characteristics.

[0002] Bibliographic details of certain publications referred to by author in this
10 specification are collected at the end of the description.

BACKGROUND OF THE INVENTION

[0003] Banana is one of the world's most important fruit crops with a world production of approximately 98 million tonnes annually (FAO, 2001). However, as with many monocultures, banana is affected by a range of fungal, viral, bacterial and
15 nematode diseases, which cause severe economical losses every year.

[0004] Fusarium wilt is one of the most destructive and notorious diseases of banana. It is also known as Panama disease, in recognition of the extensive damage it caused in export plantations in this Central American country. By 1960, Fusarium wilt had destroyed an estimated 40,000 ha of 'Gros Michel' (AAA), causing the export
20 industry to convert to cultivars in the Cavendish subgroup (AAA) (Ploetz and Pegg, 2000). Fusarium wilt is caused by the soilborne hyphomycete, *Fusarium oxysporum* Schlecht. f. sp. *cubense*. It is one of more than 120 formae speciales (special forms) of *F. oxysporum* that cause vascular wilts of flowering plants. This pathogen affects species of *Musa* and *Heliconia*, and strains have been classified into four physiological races
25 based on pathogenicity to host cultivars in the field (race 1, 'Gros Michel'; race 2, 'Bluggoe'; race 3, *Heliconia* spp.; and race 4, Cavendish cultivars and all cultivars susceptible to race 1 and 2). Until recently, race 4 had only been recorded to cause serious losses in the subtropical regions of Australia, South Africa, the Canary Islands, and Taiwan. If this race were to become established in the Americas, the world export
30 industries would be severely affected, as there is no widely accepted replacement for Cavendish cultivars (Bentley *et al.*, 1998).

[0005] In general, effective chemical control measures do not exist. In work conducted in South Africa, methyl bromide significantly reduced disease incidence, but was effective for only three years due to recolonisation of the fumigated areas by the pathogen. Studies on the biological and cultural control of this disease have begun only recently. *Arbuscular mycorrhizal* fungi have been shown to reduce disease severity in short-term green house studies, but results from long term field studies have not been reported (Ortiz *et al.*, 1995). Tissue-culture plantlets are free of pathogens and should be used to establish new plantings whenever possible. However the expense of plantlets may make their use in subsistence agriculture impractical. Genetic resistance offers the greatest opportunity for managing this disease in infested soils (Ortiz *et al.*, 1995).

[0006] Plants recognise and resist many invading pathogens by inducing a rapid defence response, termed the hypersensitive response (HR). The HR results in localised cell and tissue death at the site of infection, which constrains further spread of the infection. This local response often triggers non-specific resistance throughout the plant, a phenomenon known as systemic acquired resistance (SAR). Once triggered, SAR provides resistance to a wide range of pathogens for days. The HR and SAR depend on interaction between a dominant or semidominant resistance gene (R) product in the plant and a corresponding dominant phytopathogen avirulence gene (Avr) product (Baker *et al.*, 1997). A loss or alteration to either the plant R gene or the pathogen Avr gene leads to disease (compatibility) (Hammond-Kosack and Jones, 1997).

[0007] The R proteins provide resistance to pathogens as diverse as fungi, bacteria, viruses, nematodes and insects. Eight classes of R genes have been defined according to the structural characteristics of their predicted protein: (1) Cytoplasmic toxin reductase enzymes; (2) intracellular protein kinases; (3) receptor kinase-like protein with two tandem protein kinase domain; (4) receptor-like protein kinases with an extracellular leucine-rich repeat (LRR) domain; (5) intracellular LRR proteins with a nucleotide binding site (NBS) and leucine zipper (LZ) motif; (6) intracellular NBS-LRR proteins with a region with similarity to the Toll and interleukin-1 receptor (TIR) proteins; (7) LRR proteins that encode membrane-bound extracellular proteins; and (8) LZ proteins that encode membrane-bound intracellular proteins (Figure 1). With a few exceptions, all R genes have been cloned by a map-based cloning approach.

[0008] The NBS-LRR class is by far the largest group of resistance proteins with more than 30 cloned genes to date. Two subgroups within the NBS-LRR class have

been recognised by the presence or absence of an amino N-terminal region (TIR domain) with amino acid sequence similarity to the cytoplasmic signalling domains of the Toll and interleukin-1 receptors (Meyer *et al.*, 1999; Pan *et al.*, 2000).

[0009] The N-terminal of some NBS-LRR proteins is similar to the cytoplasmic effector domain of the *Drosophila melanogaster* and human TOLL and interleukin-1 receptors (the TIR domain) (Hammond-Kosack and Jones, 1997). Other NBS-LRR proteins have different N-terminal domains, which often contain putative leucine-zipper (LZ) motifs. Mutational analysis in *Arabidopsis* revealed that TIR-NBS-LRR and LZ-NBS-LRR proteins employ different signalling pathways. Proteins in the TIR effector domain signal via a pathway that includes EDS1, a predicted lipase, whereas most LZ-NBS-LRR proteins examined employ the membrane-associated NDR1 protein (Aarts *et al.*, 1998). There is no apparent correlation between pathogen type and the NBS-LRR subclass used by plants to detect these pathogens (Ellis and Jones 1998). All this evidence is consistent with the hypothesis of Aarts *et al.*, (1998), who suggested that there may be two downstream pathways triggered by R genes, with the structure of the R protein determining which downstream factors are required. Other recent results have shown that the situation may not be this simple. Two R genes from *Arabidopsis*, RPP8 and RPP13 (both LZ-NBS-LRR proteins), require neither EDS1 nor NDR1, suggesting that there is at least a third pathway for the transduction of R-gene signals (Glazebrook, 2001). Although many studies on different R genes have suggested that the R-protein LRR domain makes the major contribution to the unique recognition capacity of individual R genes, recent analyses of the L allelic series has shown that the TIR domain can also contribute to this capacity. Thus, it is possible that the LRR are necessary but not sufficient for the specific recognition of Avr proteins and that LRR and amino-terminal domains have co-evolved to function in a coordinate manner. (Zachary, 2001).

[0010] The central NBS domain comprises three motifs predicted to bind ATP or GTP, and several conserved motifs whose functions are not known (Hammond-Kosack and Jones, 1997). This region has homology to two activators of apoptosis in animal cells: APAF-1 and CED. By analogy to these well-characterised regulators of programmed cell death, the corresponding domain in NBS-LRR proteins might operate as an intramolecular signal transducer (Van der Biezen and Jones, 1998; Aravind *et al.*, 1999). Domain swaps involving several flax L alleles reveal a requirement for

intramolecular interactions and, thus, NBS-LRR proteins might serve as adaptor molecules that link recognition and signal delivery. For example, Avr signals perceived by the LRR might initiate nucleotide hydrolysis at the NBS domain. This might provide the energy necessary for a conformational change in the NBS-LRR protein, exposing its
5 N-terminal effector portion, to trigger a defence response (Van der Biezen and Jones, 1998).

[0011] LRR domain is thought to be involved in ligand-binding and pathogen recognition. LRR contain leucines or other hydrophobic residues at regular intervals and can also contain regularly spaced prolines and asparagines (Bent, 1996). Comparative
10 analyses of the LRR domain shows hypervariability suggesting diversification due to selection pressures. This indicates that recognition specificity resides in this part of the LRR. By analyses of *in vivo* and *in vitro* generated recombinants between different flax L alleles, Ellis *et al.* (1997) confirmed experimentally that the LRR constitute the principal determinant of specificity for Avr products. Differential specificities of R
15 proteins are often associated with duplications, deletions and sequence exchanges within the regions that encode the LRR. Recently, the LRR-like domain of the rice resistance protein Pita was shown to be required for interaction with Avr-Pita in the yeast two-hybrid system. Furthermore, mutation in either Avr-Pita Pita that abolished resistance also abolished the interaction *in vitro*. This is the first demonstrated
20 interaction between an LRR-containing R protein and its cognate Avr protein (Jia *et al.*, 2000).

[0012] Some of the resistance genes isolated to date have been transferred to susceptible cultivars of the same species or different species with successful results. For example, the N gene for resistance to Tobacco mosaic virus (TMV) has been transferred
25 to tomato and gives resistance in this species to TMV (Whitham *et al.*, 1996). The Bs2 gene, which encodes *Xanthomonas* resistance in pepper, has been cloned and transferred to tomato, a crop species in which the number of useful resistance genes to this pathogen is limited (Tai *et al.*, 1999). However, the RPS2 gene from *Arabidopsis* is non-functional in transgenic tomato and this phenomenon has been referred to as
30 'restricted taxonomic functionality' (Tai *et al.*, 1999). These data suggest that there may be difficulties in wide, cross-species resistance-gene transfer, in certain instances, due to R gene specificity (Ellis *et al.*, 2000).

[0013] The ability to isolate and transfer R genes eliminates the issue of retention of unwanted and genetically linked germoplasm, an important problem associated with classical breeding. Although disease-resistance transgenic plants are not yet available commercially, future product development seems likely as our current level of understanding of pathogenesis and plant defence improves (Stuiver and Custers 2002).

[0014] Despite the progress in R gene biology, however, no resistance genes have been isolated to date, which can confer resistance to destructive banana diseases in susceptible cultivars.

[0015] In work leading up to the present invention, four genotypes of banana were investigated to identify candidate R genes that would confer resistance to race 4 of *Fusarium oxysporum fsp cubense*. These genotypes were as follows: Cavendish, which is resistant to race 1 but susceptible to race 4; Calcutta 4, which is resistant to race 1 and race 4; three progeny of *Musa acuminata* spp *malaccensis*, which are susceptible to race 4; and three progeny of *Musa acuminata* spp *malaccensis*, which are resistant to race 4. Five families of R genes were identified from this investigation, all of which were present in the genomes of each of the genotypes but which had slightly different sequences. Surprisingly, two of these families (RGA2 and RGA5) were found to share some sequence similarity with the I2 R gene, which confers resistance to Fusarium wilt in tomatoes. In addition RGA2 was shown to be transcribed in the three resistant *Musa acuminata* spp *malaccensis* progeny but not in the three susceptible progeny. These discoveries have been reduced to practice in compositions and methods for modulating disease resistance, especially fungal resistance, in plants including banana and in plants and plant parts, especially genetically modified plants, plant cells, tissues and seeds, having modified disease resistance, as described hereafter.

SUMMARY OF THE INVENTION

[0016] Accordingly, in one aspect, the present invention provides isolated polynucleotides that confer disease resistance to a plant, especially resistance to diseases caused by fungal pathogens. These polynucleotides are generally selected from: (a) a polynucleotide comprising a nucleotide sequence that encodes a polypeptide conferring disease resistance to a plant, the sequence sharing at least 30% sequence identity with the sequence set forth in SEQ ID NO: 1 or 3, or a complement thereof; (b) a polynucleotide comprising a portion of at least 300 contiguous nucleotides of the sequence set forth in SEQ ID NO: 1 or 3, or a complement thereof, wherein the portion encodes a polypeptide that confers disease resistance to a plant; (c) a polynucleotide comprising a nucleotide sequence that encodes a polypeptide comprising the amino acid sequence set forth in SEQ ID NO: 2 or 4; (d) a polynucleotide comprising a nucleotide sequence that encodes a portion of a polypeptide comprising the amino acid sequence of SEQ ID NO: 2 or 4, wherein the portion retains the ability to confer disease resistance to a plant and comprises at least 100 contiguous amino acid residues of SEQ ID NO: 2 or 4; (e) a polynucleotide comprising a nucleotide sequence that encodes a polypeptide that shares at least 50% sequence similarity with at least a portion of at least 300 contiguous amino acid residues of the sequence set forth in SEQ ID NO: 2 or 4, wherein the polypeptide confers disease resistance to a plant; (f) a polynucleotide comprising a nucleotide sequence that encodes a polypeptide that confers disease resistance to a plant, wherein the polynucleotide hybridises to the sequence of (a), (b), (c), (d), (e) or a complement thereof, under at least low stringency conditions; and (g) a polynucleotide comprising a portion of at least 15 contiguous nucleotides of SEQ ID NO: 1 or 3, or a complement thereof, wherein the portion hybridises to a sequence of (a), (b), (c), (d), (e) or a complement thereof, under at least low stringency conditions.

[0017] In another aspect, the present invention provides nucleic acid constructs for conferring disease resistance to a plant. These constructs generally comprise a polynucleotide as broadly described operably connected to a regulatory element, which is operable in the plant. In certain embodiments, the construct is in the form of a vector, especially an expression vector.

[0018] In yet another aspect, the present invention provides isolated host cells containing a nucleic acid construct or vector as broadly described above. In certain advantageous embodiments, the host cells are plant cells. In some embodiments, the

plant cells have the nucleic acid construct or expression vector stably incorporated into their nucleome, especially their genome.

[0019] In still another aspect, the present invention provides plants containing a nucleic acid construct or expression vector as broadly described above. In certain
5 desirable embodiments, the plants have the nucleic acid construct or expression vector stably incorporated into the nucleome, especially, the genome of their cells.

[0020] In a further aspect, the present invention provides probes for interrogating nucleic acid for the presence of a disease resistance conferring polynucleotide or portion thereof. These probes generally comprise a nucleotide sequence that hybridises under at
10 least low stringency conditions to a polynucleotide as broadly described above. In some embodiments, the probes consist essentially of a nucleic acid sequence which corresponds or is complementary to at least a portion of a nucleotide sequence encoding the amino acid sequence set forth in SEQ ID NO: 2 or 4, wherein the portion is at least
15 15 nucleotides in length. In other embodiments, the probes comprise a nucleotide sequence which is capable of hybridising to at least a portion of a nucleotide sequence encoding the amino acid sequence set forth in SEQ ID NO: 2 or 4 under at least low stringency conditions, wherein the portion is at least 15 nucleotides in length. In still
other embodiment, the probes comprise a nucleotide sequence that is capable of hybridising to at least a portion of SEQ ID NO: 1 or 3 under at least low stringency
20 conditions, wherein the portion is at least 15 nucleotides in length.

[0021] Another aspect of the present invention provides methods for modulating, especially stimulating or enhancing, disease resistance in a plant. These methods generally comprise introducing an expression construct or vector as broadly described above into the nucleome of the plant and regenerating stably transformed plants. In
25 some embodiments, the expression construct or vector is introduced into regenerable plant cells so as to yield transformed plant cells, which are suitably identified and selected, and which are subsequently used for regenerating differentiated plants . Typically, a transformed plant cell line is selected from the transformed plants cells for the differentiation of a genetically modified or transgenic plant. In some embodiments,
30 the regenerable cells are regenerable dicotyledonous plant cells. In other embodiments, the regenerable cells are regenerable monocotyledonous plant cells such as regenerable graminaceous monocotyledonous plant cells and especially regenerable non-graminaceous monocotyledonous plant cells. In one example, the regenerable plant cells

are regenerable banana cells. In certain advantageous embodiments, the expression of the nucleic acid construct renders the differentiated transgenic plant disease with enhanced resistance to disease particularly diseases caused by fungal pathogens, especially soil borne fungi such as *Fusarium* species. Desirably, the nucleic acid
5 construct is transmitted through a complete cycle of the differentiated transgenic plant to its progeny so that it is expressed by the progeny plants. Thus, the invention also provides seed, other plant parts, tissue, and progeny plants derived from the differentiated transgenic plant.

[0022] In still another aspect, the invention contemplates conventional plant
10 breeding methods to transfer genetic material corresponding to a polynucleotide as broadly described above *via* crossing and backcrossing to other plants, especially plants that are susceptible to a pathogenic disease, especially a disease caused by fungal pathogens such as species of *Fusarium*. In some embodiments, the genetic material will comprise naturally-occurring DNA that corresponds to a polynucleotide as broadly
15 described above. Typically, these methods will comprise the steps of: (1) sexually crossing a plant containing that genetic material with a plant from a pathogen susceptible taxon; (2) recovering reproductive material from the progeny of the cross; and (3) growing plants with enhanced resistance to the disease from the reproductive material. In some embodiments, the methods will further comprise prior to step (1):
20 identifying a plant that is resistant to the pathogenic disease by detecting expression by the plant of a polynucleotide as broadly described above. In certain advantageous embodiments, these methods will further comprise the steps of repetitively: (a) backcrossing the disease resistant progeny with disease susceptible plants from the susceptible taxon; and (b) selecting for expression of a nucleic acid sequence
25 corresponding to a polynucleotide as broadly described above (or an associated marker gene) among the progeny of the backcross, until the desired percentage of the characteristics of the susceptible taxon are present in the progeny along with the gene or genes imparting the pathogen resistance.

[0023] In another aspect of the invention, there is provided isolated polypeptides
30 that confer disease resistance to a plant. These polypeptides are generally selected from: (i) a polypeptide comprising an amino acid sequence that shares at least 50% sequence similarity with the sequence set forth in SEQ ID NO: 2 or 4; (ii) a polypeptide comprising a portion of at least 100 contiguous amino acid residues of the sequence set

forth in SEQ ID NO: 2 or 4, wherein the polypeptide confers disease resistance to a plant; (iii) a polypeptide comprising an amino acid sequence that shares at least 30% similarity with at least a portion of 100 contiguous amino acid residues of the sequence set forth in SEQ ID NO: 2 or 4; and (iv) a polypeptide comprising a portion of at least 5
5 contiguous amino acid residues of the sequence set forth in SEQ ID NO: 2 or 4, wherein the portion is immuno-interactive with an antigen-binding molecule that is immuno-interactive with a sequence of (i), (ii) or (iii).

[0024] In some embodiments, the polypeptide includes one or more and in some cases all of the following domains (the numbering refers to the consensus numbering in
10 Figure 2):

a domain which corresponds to residues 1-167 of Figure 2. This domain may be structurally similar to a coiled coil. In some embodiments, this domain can have at least 60, 70, 80, 90, 95, or 98 % sequence similarity with, or have at least 30, 40, 50, 60, 70 or 80 % sequence identity to, or differ at no more than 1, 2, 3, 4, 5, 10, 15, 20,
15 25, 30 or 40 amino acid residues from, the corresponding domain of any of the sequences presented in Figure 2;

a domain which corresponds to residues 168-536 of Figure 2. This domain may be functionally analogous to a nuclear-binding site (NBS) domain. In some embodiments, this domain can have at least 70, 80, 90, 95, or 98 % sequence similarity
20 with, or have at least 50, 60, 70, 80 or 90 % sequence identity to, or differ at no more than 1, 2, 3, 4, 5, 10, 15, 20, 25, 30 or 40 amino acid residues from, the corresponding domain of any of the sequences presented in Figure 2; and

a domain which corresponds to residues 537-1476 of Figure 2. This domain may be functionally analogous to a leucine-rich repeat (LRR) domain. In some
25 embodiments, this domain can have at least 60, 70, 80, 90, 95, or 98 % sequence similarity with, or have at least 30, 40, 50, 60, 70, 80 or 90 % sequence identity to, or differ at no more than 1, 2, 3, 4, 5, 10, 15, 20, 25, 30 or 40 amino acid residues from, the corresponding domain of any of the sequences presented in Figure 2.

[0025] In some embodiments, it may be desirable to conserve one or more of the
30 residues in the above regions, which are conserved between the sequences presented in Figure 2, wherein the conserved amino acid residues correspond to identical residues or to residues belonging to the same class or subclass of amino acid residues.

[0026] In some embodiments, the domain corresponding to residues 1-167 of Figure 2 comprises a sequence according to Formula (I):

5 Ser- Φ_{aa1} - Φ_{aa2} Zaa-Xaa₁- Φ_{aa3} - Φ_{aa4} Xaa₂-Baa₁ Σ_{aa1} -Xaa₃-Asn-Xaa₄-Xaa₅- Φ_{aa5} -
Xaa₆-Xaa₇-Leu-Xaa₈-Xaa₉-Xaa₁₀-Xaa₁₁-Xaa₁₂-Xaa₁₃-Baa₂-Xaa₁₄- $\mathring{A}aa_1$ -Leu-Xaa₁₅-
Xaa₁₆-Leu-Xaa₁₇-Xaa₁₈- Σ_{aa2} -Leu-Leu-Arg-Xaa₁₉-His- Σ_{aa3} - Φ_{aa6} -Leu- $\mathring{A}aa_2$ - Ω_{aa1} -
Ala- Ω_{aa2} - Σ_{aa4} -Arg-Xaa₂₀-Xaa₂₁-Xaa₂₂-Xaa₂₃-Xaa₂₄-Xaa₂₅-Xaa₂₆-Ser-Leu-Val-
Xaa₂₇- Φ_{aa7} - Φ_{aa8} -Xaa₂₈-Xaa₂₉-Leu-Lys- $\mathring{A}aa_3$ -Xaa₃₀-Ala-Tyr-Asp-Ala- $\mathring{A}aa_4$ -Asp-
15 Φ_{aa9} -Leu- $\mathring{A}aa_5$ -Glu- Φ_{aa10} -Glu-Xaa₃₁-Xaa₃₂-Ala-Xaa₃₃-Baa₃-Xaa₃₄-Lys-Val (I)

wherein: each of Φ_{1-10} is a hydrophobic amino acid residue,

10 Zaa is a neutral/polar amino acid residue,

each of Σ_{aa1-4} is a small amino acid residue,

each of Baa₁₋₃ is a basic amino acid residue,

each of $\mathring{A}aa_{1-5}$ is an acidic amino acid residue,

each of Ω_{aa1-2} is a charged amino acid residue, and

15 Xaa₁₋₃₃ are each independently selected from any amino acid residue.

[0027] In some embodiments, Zaa is selected from Gln or Asn.

[0028] In some embodiments, Φ_{aa1} is selected from Phe or Leu. In some
embodiments, Φ_{aa2} is selected from Ile or Val. In some embodiments, Φ_{aa3} is selected
from Leu or Ile. In some embodiments, Φ_{aa4} is selected from Leu or Phe. In some
20 embodiments, Φ_{aa5} is selected from Ile or Val. In some embodiments, Φ_{aa6} is selected
from Ile or Leu. In some embodiments, Φ_{aa7} is selected from Leu or Trp. In some
embodiments, Φ_{aa8} is selected from Val or Leu. In some embodiments, Φ_{aa9} is selected
from Leu or Ile. In some embodiments, Φ_{aa10} is selected from Leu or Trp.

[0029] In some embodiments, Σ_{aa1} is selected from Ala Ser. In some embodiments,
25 Σ_{aa2} is selected from Ser or Thr. In some embodiments, Σ_{aa3} is selected from Ala Ser.
In some embodiments, Σ_{aa4} is selected from Thr or Ala.

[0030] In some embodiments, Baa₁ is selected from Lys or Arg. In some
embodiments, Baa₂ is selected from His or Arg. In some embodiments, Baa₃ is selected
from Lys or Arg.

30 [0031] In some embodiments, each of $\mathring{A}aa_{1-5}$ is independently selected from Asp or
Glu.

[0032] In some embodiments, Ωaa_1 is selected from Lys or Glu. In some embodiments, Ωaa_2 is selected from Glu or Lys.

[0033] In some embodiments Xaa_1 is a small or acidic amino acid residue, e.g., Xaa_1 is selected from Thr or Glu. In some embodiments, Xaa_2 is an acidic or neutral/polar amino acid residue, e.g., Xaa_2 is selected from Asp or Asn. In some embodiments, Xaa_3 is a small or hydrophobic amino acid residue, e.g., Xaa_3 is selected from Ser or Ile. In some embodiments, Xaa_4 is a neutral/polar or hydrophobic amino acid residue, e.g., Xaa_4 is selected from Cys or Leu. In some embodiments, Xaa_5 is a small or hydrophobic amino acid residue, e.g., Xaa_5 is selected from Ala Ile. In some embodiments, Xaa_6 is a neutral/polar or small amino acid residue, e.g., Xaa_6 is selected from Gln or Ala. In some embodiments, Xaa_7 is a neutral/polar or acidic amino acid residue, e.g., Xaa_7 is selected from Gln or Glu. In some embodiments, Xaa_8 is a small or basic amino acid residue, e.g., Xaa_8 is selected from Ala Arg. In some embodiments, Xaa_9 is a basic or hydrophobic amino acid residue, e.g., Xaa_9 is selected from Arg or Leu.

[0034] In some embodiments, Xaa_{10} is a basic or neutral/polar amino acid residue, e.g., Xaa_{10} is selected from Arg or Gln. In some embodiments, Xaa_{11} is a basic or hydrophobic amino acid residue, e.g., Xaa_{11} is selected from Arg or Leu. In some embodiments, Xaa_{12} is a small or neutral/polar amino acid residue, e.g., Xaa_{12} is selected from Arg or Gln. In some embodiments, Xaa_{13} is a hydrophobic or small amino acid residue, e.g., Xaa_{13} is selected from Leu or Ala. In some embodiments, Xaa_{14} is an acid or small amino acid residue, e.g., Xaa_{14} is selected from Asp or Ala. In some embodiments, Xaa_{15} is a basic or neutral/polar amino acid residue, e.g., Xaa_{15} is selected from Arg or Asn. In some embodiments, Xaa_{16} is a basic or neutral/polar amino acid residue, e.g., Xaa_{16} is selected from Arg or Asn. In some embodiments, Xaa_{17} is a basic or neutral/polar amino acid residue, e.g., Xaa_{17} is selected from Arg or Gln. In some embodiments, Xaa_{18} is a small or basic amino acid residue, e.g., Xaa_{18} is selected from Thr or Arg. In some embodiments, Xaa_{19} is a hydrophobic or small amino acid residue, e.g., Xaa_{19} is selected from Ile or Thr.

[0035] In some embodiments, Xaa_{20} is a hydrophobic or basic amino acid residue, e.g., Xaa_{20} is selected from Trp or Arg. In some embodiments, Xaa_{21} is absent or is a neutral/polar amino acid residue, e.g., Asn. In some embodiments, Xaa_{22} is a basic or hydrophobic amino acid residue, e.g., Xaa_{22} is selected from His or Met. In some

embodiments, Xaa₂₃ is a basic or small amino acid residue, e.g., Xaa₂₃ is selected from Lys or Thr. In some embodiments, Xaa₂₄ is a neutral/polar or acidic amino acid residue, e.g., Xaa₂₄ is selected from Asn or Asp. In some embodiments, Xaa₂₅ is a small or basic amino acid residue, e.g., Xaa₂₅ is selected from Thr or Lys. In some embodiments, Xaa₂₆ is an acidic or hydrophobic amino acid residue, e.g., Xaa₂₆ is selected from Glu or Leu. In some embodiments, X₂₇ is a basic or hydrophobic or amino acid residue, e.g., Xaa₂₇ is selected from Arg or Met. In some embodiments, Xaa₂₈ is a neutral/polar or acidic amino acid residue, e.g., Xaa₂₈ is selected from Gln or Glu. In some embodiments, Xaa₂₉ is a small or hydrophobic amino acid residue, e.g., Xaa₂₉ is selected from Ala Trp.

[0036] In some embodiments, Xaa₃₀ is a hydrophobic or small amino acid residue, e.g., Xaa₃₀ is selected from Tyr or Ala. In some embodiments, Xaa₃₁ is a neutral/polar or small amino acid residue, e.g., Xaa₃₁ is selected from Gln or Ala. In some embodiments, Xaa₃₂ is a small or hydrophobic amino acid residue, e.g., Xaa₃₂ is selected from Ala Ile. In some embodiments, Xaa₃₃ is a neutral/polar or hydrophobic amino acid residue, e.g., Xaa₃₃ is selected from Gln or Leu.

[0037] In some embodiments, the domain corresponding to residues 168-536 of Figure 2 comprises a sequence according to Formula (II):

Arg-Xaa₁-Xaa₂-Thr-Σaa₁-Ser-Φaa₁-Leu-Thr-Glu-Σaa₂-Xaa₃-Φaa₂-Φaa₃-Gly-Arg-
 Xaa₄-Gln-Åaa₁-Baa₁-Glu-Xaa₅-Φaa₄-Φaa₅-Ωaa₁-Leu-Leu-Leu-Åaa₂-Σaa₃-Σaa₄-
 Xaa₆-Gly-Xaa₇-Xaa₈-Σaa₅-Phe-Σaa₆-Val-Φaa₆-Pro-Φaa₇-Val-Gly-Φaa₈-Gly-Gly-
 Xaa₉-Gly-Lys-Thr-Thr-Leu-Σaa₇-Gln-Leu-Φaa₉-Φaa₁₀-Asn-Asp-Xaa₁₀-Arg-Val-
 Xaa₁₁-Xaa₁₂-Xaa₁₃-Phe-Xaa₁₄-Leu-Baa₂-Φaa₁₁-Trp-Val-Cys-Val-Ser-Asp-Xaa₁₅-
 Phe-Xaa₁₆-Val-Lys-Arg-Φaa₁₂-Thr-Baa₃-Glu-Ile-Xaa₁₇-Glu-Xaa₁₈-Ala-Thr-Xaa₁₉-
 Xaa₂₀-Ωaa₂-Xaa₂₁-Xaa₂₂-Asp-Xaa₂₃-Xaa₂₄-Asn-Leu-Xaa₂₅-Xaa₂₆-Leu-Gln-Xaa₂₇-
 Xaa₂₈-Leu-Lys-Glu-Ωaa₃-Ile-Xaa₂₉-Σaa₈-Xaa₃₀-Xaa₃₁-Phe-Leu-Leu-Val-Leu-Asp-
 Asp-Val-Trp-Xaa₃₂-Glu-Xaa₃₃-Xaa₃₄-Xaa₃₅-Ωaa₄-Trp-Glu-Xaa₃₆-Leu-Xaa₃₇-Ala-
 Pro-Leu-Ωaa₅-Xaa₃₈-Σaa₉-Σaa₁₀-Arg-Gly-Ser-Xaa₃₉-Val-Ile-Val-Thr-Thr-Xaa₄₀-
 Xaa₄₁-Xaa₄₂-Lys-Φaa₁₃-Ala-Xaa₄₃-Φaa₁₄-Xaa₄₄-Gly-Thr-Met-Ωaa₆-Xaa₄₅-Φaa₁₅-
 Xaa₄₆-Leu-Åaa₃-Xaa₄₇-Leu-Xaa₄₈-Åaa₄-Asp-Xaa₄₉-Xaa₅₀-Trp-Xaa₅₁-Leu-Φaa₁₆-
 Ωaa₇-Xaa₅₂-Xaa₅₃-Σaa₁₁-Phe-Xaa₅₄-Xaa₅₅-Xaa₅₆-Xaa₅₇-Xaa₅₈-Σaa₁₂-Xaa₅₉-Xaa₆₀-
 Xaa₆₁-Xaa₆₂-Ωaa₈-Φaa₁₇-Glu-Xaa₆₃-Ile-Gly-Arg-Lys-Ile-Ala-Xaa₆₄-Lys-Φaa₁₈-

Xaa₆₅-Gly-Xaa₆₆-Pro-Φaa₁₉-Σaa₁₃-Ala-Xaa₆₇-Σaa₁₄-Φaa₂₀-Gly-Xaa₆₈-Φaa₂₁-Leu-
 Arg-Xaa₆₉-Ωaa₉-Xaa₇₀-Σaa₁₅-Xaa₇₁-Xaa₇₂-Xaa₇₃-Trp-Arg-Xaa₇₄-Φaa₂₂-Φaa₂₃-
 Glu-Σaa₁₆-Glu-Xaa₇₅-Trp-Xaa₇₆-Φaa₂₄-Pro-Xaa₇₇-Ala-Xaa₇₈-Xaa₇₉-Åaa₅-Φaa₂₅-
 Leu-Σaa₁₇-Xaa₈₀-Leu-Xaa₈₁-Xaa₈₂-Ser-Tyr-Xaa₈₃-Xaa₈₄-Leu-Pro-Σaa₁₈-Xaa₈₅-
 5 Leu-Baa₄-Xaa₈₆-Cys-Phe-Ala-Phe-Cys-Ala-Φaa₂₆-Phe-Xaa₈₇-Lys-Xaa₈₈-Tyr-
 Xaa₈₉-Phe-Xaa₉₀-Lys-Ωaa₁₀-Xaa₉₁-Leu-Ile-Xaa₉₂-Xaa₉₃-Trp-Ile-Ala-Xaa₉₄-Xaa₉₅-
 Φaa₂₇-Ile

wherein: each of Φ₁₋₂₇ is a hydrophobic amino acid residue,
 each of Σaa₁₋₁₈ is a small amino acid residue,
 10 each of Baa₁₋₄ is a basic amino acid residue,
 each of Åaa₁₋₅ is an acidic amino acid residue,
 each of Ωaa₁₋₁₀ is a charged amino acid residue, and
 Xaa₁₋₉₅ are each independently selected from any amino acid residue.

[0038] In some embodiments, Σaa₁ is selected from Ser or Thr. In some
 15 embodiments, Σaa₂ is selected from Thr or Ser. In some embodiments, Σaa₃ is selected
 from Ser or Pro. In some embodiments, Σaa₄ is selected from Gly or Ser. In some
 embodiments, Σaa₅ is selected from Ser or Ala. In some embodiments, Σaa₆ is selected
 from Ser or Pro. In some embodiments, Σaa₇ is selected from Ala or Ser. In some
 embodiments, Σaa₈ is selected from Ser or Gly. In some embodiments, Σaa₉ is selected
 20 from Ala or Gly.

[0039] In some embodiments, Σaa₁₀ is selected from Ala or Gly. In some
 embodiments, Σaa₁₁ is selected from Ala or Ser. In some embodiments, Σaa₁₂ is selected
 from Pro or Ser. In some embodiments, Σaa₁₃ is selected from Ala or Gly. In some
 embodiments, Σaa₁₄ is selected from Thr or Ala. In some embodiments, Σaa₁₅ is
 25 selected from Ser or Gly. In some embodiments, Σaa₁₆ is selected from Ser or Thr. In
 some embodiments, Σaa₁₇ is selected from Pro or Ser. In some embodiments, Σaa₁₈ is
 selected from Gly or Pro.

[0040] In some embodiments, Φaa₁ is selected from Phe or Leu. In some
 embodiments, Φaa₂ is selected from Val or Ile. In some embodiments, Φaa₃ is selected
 30 from Phe or Val. In some embodiments, Φaa₄ is selected from Val or Leu. In some
 embodiments, Φaa₅ is selected from Val or Ile. In some embodiments, Φaa₆ is selected

from Leu or Val. In some embodiments, Φ_{aa7} is selected from Leu or Ile. In some embodiments, Φ_{aa8} is selected from Ile or Val. In some embodiments, Φ_{aa9} is selected from Val or Ile.

[0041] In some embodiments, Φ_{aa10} is selected from Tyr or Phe. In some
5 embodiments, Φ_{aa11} is selected from Val or Met. In some embodiments, Φ_{aa12} is
selected from Leu or Ile. In some embodiments, Φ_{aa13} is selected from Ile or Val. In
some embodiments, Φ_{aa14} is selected from Ile or Val. In some embodiments, Φ_{aa15} is
selected from Ile or Tyr. In some embodiments, Φ_{aa16} is selected from Phe or Ile. In
some embodiments, Φ_{aa17} is selected from Leu or Met. In some embodiments, Φ_{aa18} is
10 selected from Leu or Ile. In some embodiments, Φ_{aa19} is selected from Leu or Tyr.

[0042] In some embodiments, Φ_{aa20} is selected from Leu or Met. In some
embodiments, Φ_{aa21} is selected from Leu or Tyr. In some embodiments, Φ_{aa22} is
selected from Ile or Val. In some embodiments, Φ_{aa23} is selected from Met or Leu. In
some embodiments, Φ_{aa24} is selected from Leu or Met. In some embodiments, Φ_{aa25} is
15 selected from Ile or Val. In some embodiments, Φ_{aa26} is selected from Val or Leu. In
some embodiments, Φ_{aa27} is selected from Phe or Leu.

[0043] In some embodiments, B_{aa1-4} are each independently selected from Arg or
Lys.

[0044] In some embodiments, each of A_{aa1-5} is independently selected from Asp or
20 Glu.

[0045] In some embodiments, Ω_{aa1} is selected from Glu or Arg. In some
embodiments, Ω_{aa2} is selected from Glu or Arg. In some embodiments, Ω_{aa3} is selected
from Lys or Glu. In some embodiments, Ω_{aa4} is selected from Asp or Lys. In some
embodiments, Ω_{aa5} is selected from Arg or Asp. In some embodiments, Ω_{aa6} is selected
25 from Lys or Glu. In some embodiments, Ω_{aa7} is selected from Lys or Glu. In some
embodiments, Ω_{aa8} is selected from Glu or Arg. In some embodiments, Ω_{aa9} is selected
from Asp or Lys. In some embodiments, Ω_{aa10} is selected from His or Asp.

[0046] In some embodiments X_{aa1} is a basic or small amino acid residue, e.g., X_{aa1}
is selected from Arg or Gly. In some embodiments, X_{aa2} is an acidic or hydrophobic
30 amino acid residue, e.g., X_{aa2} is selected from Glu or Val. In some embodiments, X_{aa3}
is a hydrophobic or neutral/polar amino acid residue, e.g., X_{aa3} is selected from Val or

Cys. In some embodiments, Xaa₄ is an acidic or small amino acid residue, e.g., Xaa₄ is selected from Asp or Ala. In some embodiments, Xaa₅ is a basic or neutral/polar amino acid residue, e.g., Xaa₅ is selected from Lys or Asn. In some embodiments, Xaa₆ is a small or acidic amino acid residue, e.g., Xaa₆ is selected from Ser or Asp. In some
5 embodiments, Xaa₇ is absent or is a neutral/polar amino acid residue, e.g., Asn. In some embodiments, Xaa₈ is absent or is a small amino acid residue, e.g., Ser. In some embodiments, Xaa₉ is a hydrophobic or small amino acid residue, e.g., Xaa₉ is selected from Val or Ala.

[0047] In some embodiments, Xaa₁₀ is a neutral/polar or basic amino acid residue,
10 e.g., Xaa₁₀ is selected from Asn or Lys. In some embodiments, Xaa₁₁ is a small or acidic amino acid residue, e.g., Xaa₁₁ is selected from Gly or Glu. In some embodiments, Xaa₁₂ is a neutral/polar or acidic amino acid residue, e.g., Xaa₁₂ is selected from Asn or Glu. In some embodiments, Xaa₁₃ is a hydrophobic or basic amino acid residue, e.g., Xaa₁₃ is selected from Tyr or His. In some embodiments, Xaa₁₄ is a basic or small
15 amino acid residue, e.g., Xaa₁₄ is selected from His or Pro. In some embodiments, Xaa₁₅ is a neutral/polar or acidic amino acid residue, e.g., Xaa₁₅ is selected from Asn or Asp. In some embodiments, Xaa₁₆ is a neutral/polar or acidic amino acid residue, e.g., Xaa₁₆ is selected from Asn or Asp. In some embodiments, Xaa₁₇ is a hydrophobic or small amino acid residue, e.g., Xaa₁₇ is selected from Ile or Thr. In some embodiments, Xaa₁₈
20 is a small or hydrophobic amino acid residue, e.g., Xaa₁₈ is selected from Ser or Tyr. In some embodiments, Xaa₁₉ is a basic or neutral/polar amino acid residue, e.g., Xaa₁₉ is selected from Lys or Asn.

[0048] In some embodiments, Xaa₂₀ is a hydrophobic or small amino acid residue, e.g., Xaa₂₀ is selected from Val or Gly. In some embodiments, Xaa₂₁ is a neutral/polar
25 or hydrophobic amino acid residue, e.g., Xaa₂₁ is selected from Gln or Phe. In some embodiments, Xaa₂₂ is a small or hydrophobic amino acid residue, e.g., Xaa₂₂ is selected from Ser or Met. In some embodiments, Xaa₂₃ is a basic or hydrophobic amino acid residue, e.g., Xaa₂₃ is selected from Lys or Leu. In some embodiments, Xaa₂₄ is a hydrophobic or small amino acid residue, e.g., Xaa₂₄ is selected from Leu or Thr. In
30 some embodiments, Xaa₂₅ is an acidic or neutral/polar amino acid residue, e.g., Xaa₂₅ is selected from Asp or Asn. In some embodiments, Xaa₂₆ is a small or hydrophobic amino acid residue, e.g., Xaa₂₆ is selected from Thr or Met. In some embodiments, Xaa₂₇ is a neutral/polar or hydrophobic amino acid residue, e.g., Xaa₂₇ is selected from

Gln or Val. In some embodiments, Xaa₂₈ is a hydrophobic or neutral/polar amino acid residue, e.g., Xaa₂₈ is selected from Ile or Asn. In some embodiments, Xaa₂₉ is a small or basic amino acid residue, e.g., Xaa₂₉ is selected from Ala or Arg.

[0049] In some embodiments, Xaa₃₀ is an acidic or small amino acid residue, e.g.,
5 Xaa₃₀ is selected from Glu or Thr. In some embodiments, X₃₁ is a basic or small amino acid residue, e.g., Xaa₃₁ is selected from Arg or Thr. In some embodiments, Xaa₃₂ is a small or neutral/polar amino acid residue, e.g., Xaa₃₂ is selected from Ser or Asn. In some embodiments, Xaa₃₃ is a neutral/polar or acidic amino acid residue, e.g., Xaa₃₃ is selected from Asn or Asp. In some embodiments, Xaa₃₄ is a basic or small amino acid
10 residue, e.g., Xaa₃₄ is selected from Arg or Pro. In some embodiments, Xaa₃₅ is an acidic or hydrophobic amino acid residue, e.g., Xaa₃₅ is selected from Asp or Val. In some embodiments, Xaa₃₆ is a basic or small amino acid residue, e.g., Xaa₃₆ is selected from Arg or Ser. In some embodiments, Xaa₃₇ is a neutral/polar or hydrophobic amino acid residue, e.g., Xaa₃₇ is selected from Cys or Leu. In some embodiments, Xaa₃₈ is a
15 hydrophobic or small amino acid residue, e.g., Xaa₃₈ is selected from Phe or Ala. In some embodiments, Xaa₃₉ is a basic or hydrophobic amino acid residue, e.g., Xaa₃₉ is selected from Lys or Val.

[0050] In some embodiments, Xaa₄₀ is a basic or neutral/polar amino acid residue, e.g., Xaa₄₀ is selected from Arg or Gln. In some embodiments, X₄₁ is an acidic or small
20 amino acid residue, e.g., Xaa₄₁ is selected from Asp or Ser. In some embodiments, Xaa₄₂ is a small or basic amino acid residue, e.g., Xaa₄₂ is selected from Thr or Lys. In some embodiments, Xaa₄₃ is a small or acidic amino acid residue, e.g., Xaa₄₃ is selected from Ser or Asp. In some embodiments, Xaa₄₄ is a hydrophobic or small amino acid residue, e.g., Xaa₄₄ is selected from Ile or Thr. In some embodiments, Xaa₄₅ is an acidic
25 or small amino acid residue, e.g., Xaa₄₅ is selected from Glu or Pro. In some embodiments, Xaa₄₆ is a small or hydrophobic amino acid residue, e.g., Xaa₄₆ is selected from Ser or Val. In some embodiments, Xaa₄₇ is a small or acidic amino acid residue, e.g., Xaa₄₇ is selected from Gly or Glu. In some embodiments, Xaa₄₈ is a neutral/polar or small amino acid residue, e.g., Xaa₄₈ is selected from Gln or Thr. In
30 some embodiments, Xaa₄₉ is a small or acidic amino acid residue, e.g., Xaa₄₉ is selected from Ala or Asp.

[0051] In some embodiments, Xaa₅₀ is a hydrophobic or small amino acid residue, e.g., Xaa₅₀ is selected from Tyr or Ser. In some embodiments, X₅₁ is an acidic or small

amino acid residue, e.g., Xaa₅₁ is selected from Glu or Ser. In some embodiments, Xaa₅₂ is a basic or small amino acid residue, e.g., Xaa₅₂ is selected from Lys or Ser. In some embodiments, Xaa₅₃ is a neutral/polar or basic amino acid residue, e.g., Xaa₅₃ is selected from Cys or His. In some embodiments, Xaa₅₄ is a small or basic amino acid residue, e.g., Xaa₅₄ is selected from Gly or Arg. In some embodiments, Xaa₅₅ is a small or acidic amino acid residue, e.g., Xaa₅₅ is selected from Ser or Glu. In some embodiments, Xaa₅₆ is a hydrophobic or small amino acid residue, e.g., Xaa₅₆ is selected from Val or Ala. In some embodiments, Xaa₅₇ is a neutral/polar or small amino acid residue, e.g., Xaa₅₇ is selected from Asn or Ser. In some embodiments, Xaa₅₈ is absent or is a neutral/polar amino acid residue, e.g., Cys. In some embodiments, Xaa₅₉ is a neutral/polar or small amino acid residue, e.g., Xaa₅₉ is selected from Gln or Ser.

[0052] In some embodiments, Xaa₆₀ is an acidic or small amino acid residue, e.g., Xaa₆₀ is selected from Glu or Pro. In some embodiments, Xaa₆₁ is a basic or neutral/polar amino acid residue, e.g., Xaa₆₁ is selected from His or Asn. In some embodiments, Xaa₆₂ is a hydrophobic or small amino acid residue, e.g., Xaa₆₂ is selected from Leu or Pro. In some embodiments, Xaa₆₃ is a hydrophobic or acidic amino acid residue, e.g., Xaa₆₃ is selected from Val or Glu. In some embodiments, Xaa₆₄ is a small or basic amino acid residue, e.g., Xaa₆₄ is selected from Gly or Lys. In some embodiments, Xaa₆₅ is a basic or small amino acid residue, e.g., Xaa₆₅ is selected from Lys or Ser. In some embodiments, Xaa₆₆ is a small or hydrophobic amino acid residue, e.g., Xaa₆₆ is selected from Ser or Leu. In some embodiments, Xaa₆₇ is a basic or small amino acid residue, e.g., Xaa₆₇ is selected from Lys or Thr. In some embodiments, Xaa₆₈ is a small or basic amino acid residue, e.g., Xaa₆₈ is selected from Ser or Arg. In some embodiments, Xaa₆₉ is a hydrophobic or small amino acid residue, e.g., Xaa₆₉ is selected from Leu or Ser.

[0053] In some embodiments, Xaa₇₀ is a hydrophobic or basic amino acid residue, e.g., Xaa₇₀ is selected from Val or His. In some embodiments, Xaa₇₁ is a neutral/polar or acidic amino acid residue, e.g., Xaa₇₁ is selected from Gln or Glu. In some embodiments, Xaa₇₂ is an acidic or small amino acid residue, e.g., Xaa₇₂ is selected from Glu or Ser. In some embodiments, Xaa₇₃ is a basic or small amino acid residue, e.g., Xaa₇₃ is selected from His or Ser. In some embodiments, Xaa₇₄ is a small or acidic amino acid residue, e.g., Xaa₇₄ is selected from Thr or Glu. In some embodiments, Xaa₇₅ is a hydrophobic or small amino acid residue, e.g., Xaa₇₅ is selected from Val or

Thr. In some embodiments, Xaa₇₆ is a neutral/polar or acidic amino acid residue, e.g., Xaa₇₆ is selected from Gln or Glu. In some embodiments, Xaa₇₇ is a neutral/polar or small amino acid residue, e.g., Xaa₇₇ is selected from Gln or Pro. In some embodiments, Xaa₇₈ is an acidic or small amino acid residue, e.g., Xaa₇₈ is selected from Glu or Ala.

- 5 In some embodiments, Xaa₇₉ is a neutral/polar or small amino acid residue, e.g., Xaa₇₉ is selected from Asn or Ser.

[0054] In some embodiments, Xaa₈₀ is a hydrophobic or small amino acid residue, e.g., Xaa₈₀ is selected from Val or Ala. In some embodiments, Xaa₈₁ is a hydrophobic or basic amino acid residue, e.g., Xaa₈₁ is selected from Trp or Arg. In some embodiments,

- 10 Xaa₈₂ is a hydrophobic or basic amino acid residue, e.g., Xaa₈₂ is selected from Leu or Arg. In some embodiments, Xaa₈₃ is a neutral/polar or acidic amino acid residue, e.g., Xaa₈₃ is selected from Gln or Asp. In some embodiments, Xaa₈₄ is a basic or neutral/polar amino acid residue, e.g., Xaa₈₄ is selected from His or Asn. In some embodiments, Xaa₈₅ is a basic or neutral/polar amino acid residue, e.g., Xaa₈₅ is selected from His or Gln. In some embodiments, Xaa₈₆ is a neutral/polar or hydrophobic amino acid residue, e.g., Xaa₈₆ is selected from Gln or Leu. In some embodiments, Xaa₈₇ is a basic or small amino acid residue, e.g., Xaa₈₇ is selected from His or Thr. In some embodiments, Xaa₈₈ is an acidic or small amino acid residue, e.g., Xaa₈₈ is selected from Asp or Gly. In some embodiments, Xaa₈₉ is a hydrophobic or basic amino acid residue, e.g., Xaa₈₉ is selected from Leu or Arg.

[0055] In some embodiments, Xaa₉₀ is a hydrophobic or basic amino acid residue, e.g., Xaa₉₀ is selected from Tyr or Arg. In some embodiments, Xaa₉₁ is an acidic or small amino acid residue, e.g., Xaa₉₁ is selected from Glu or Thr. In some embodiments, Xaa₉₂ is a neutral/polar or basic amino acid residue, e.g., Xaa₉₂ is selected from Gln or His. In some embodiments, Xaa₉₃ is a small or hydrophobic amino acid residue, e.g., Xaa₉₃ is selected from Thr or Met. In some embodiments, Xaa₉₄ is an acidic or neutral/polar amino acid residue, e.g., Xaa₉₄ is selected from Glu or Gln. In some embodiments, Xaa₉₅ is a small or neutral/polar amino acid residue, e.g., Xaa₉₅ is selected from Gly or Asn.

- 30 [0056] In some embodiments, the domain corresponding to residues 537-1476 of Figure 2 comprises a sequence according to Formula (II):

Leu-Xaa₁-Ωaa₁-Xaa₂-Φaa₁-Phe-Baa₁-Xaa₃-Leu-Xaa₄-Arg-Ile-Baa₂-Val-Leu-Xaa₅-
Φaa₂-Xaa₆-Xaa₇-Cys-Xaa₈-Φaa₃-Baa₃-Xaa₉-Leu-Pro-Xaa₁₀-Xaa₁₁-Φaa₄-Gly-

Xaa₁₂-Leu-Xaa₁₃-Xaa₁₄-Leu-Arg-Tyr-Leu-Xaa₁₅-Φaa₅-Ser-Xaa₁₆-Asn-Σaa₁-Xaa₁₇-
 Ile-Gln-Arg-Leu-Pro-Glu-Ser-Φaa₆-Xaa₁₈-Ωaa₂-Leu-Xaa₁₉-Xaa₂₀-Leu-Gln-Σaa₂-
 Leu-Xaa₂₁-Leu-Xaa₂₂-Gly-Cys-Xaa₂₃-Leu-Xaa₂₄-Xaa₂₅-Φaa₇-Pro-Xaa₂₆-Σaa₃-
 Met-Ser-Baa₄-Leu-Φaa₈-Xaa₂₇-Leu-Arg-Gln-Leu-Baa₅-Xaa₂₈-Xaa₂₉-Xaa₃₀-Åaa₁-
 5 Φaa₉-Ile-Σaa₄-Ωaa₃-Ile-Xaa₃₁-Ωaa₄-Val-Gly-Baa₆-Leu-Ile-Xaa₃₂-Leu-Gln-Glu-
 Leu-Xaa₃₃-Ala-Φaa₁₀-Xaa₃₄-Val-Xaa₃₅-Xaa₃₆-Baa₇-Xaa₃₇-Gly-Xaa₃₈-Xaa₃₉-Φaa₁₁-
 Ala-Glu-Leu-Ser-Σaa₅-Φaa₁₂-Xaa₄₀-Gln-Leu-Baa₈-Σaa₆-Xaa₄₁-Leu-Xaa₄₂-Ile-
 Xaa₄₃-Asn-Leu-Xaa₄₄-Asn-Val-Xaa₄₅-Xaa₄₆-Xaa₄₇-Ωaa₅-Glu-Σaa₇-Xaa₄₈-Lys-
 Ala-Baa₉-Leu-Ωaa₆-Ωaa₇-Lys-Gln-Xaa₄₉-Leu-Ωaa₈-Xaa₅₀-Leu-Åaa₂-Leu-Ωaa₉-
 10 Trp-Ala-Xaa₅₁-Gly-Xaa₅₂-Xaa₅₃-Xaa₅₄-Xaa₅₅-Xaa₅₆-Xaa₅₇-Xaa₅₈-Glu-Xaa₅₉-
 Xaa₆₀-Xaa₆₁-Xaa₆₂-Ωaa₁₀-Ωaa₁₁-Val-Leu-Xaa₆₃-Gly-Leu-Xaa₆₄-Pro-His-Xaa₆₅-
 Xaa₆₆-Leu-Baa₁₀-Xaa₆₇-Leu-Σaa₈-Ile-Baa₁₁-Xaa₆₈-Tyr-Σaa₉-Gly-Σaa₁₀-Σaa₁₁-
 Xaa₆₉-Pro-Ser-Trp-Φaa₁₃-Xaa₇₀-Xaa₇₁-Xaa₇₂-Φaa₁₄-Leu-Pro-Asn-Φaa₁₅-Xaa₇₃-
 Thr-Φaa₁₆-Baa₁₂-Leu-Ωaa₁₂-Xaa₇₄-Cys-Σaa₁₂-Arg-Leu-Xaa₇₅-Xaa₇₆-Leu-Σaa₁₃-
 15 Xaa₇₇-Φaa₁₇-Gly-Gln-Leu-Xaa₇₈-Xaa₇₉-Leu-Baa₁₃-Xaa₈₀-Leu-His-Φaa₁₈-Ωaa₁₃-
 Xaa₈₁-Met-Σaa₁₄-Xaa₈₂-Val-Baa₁₄-Gln-Φaa₁₉-Xaa₈₃-Xaa₈₄-Xaa₈₅-Φaa₂₀-Xaa₈₆-
 Gly-Xaa₈₇-Σaa₁₅-Ωaa₁₄-Xaa₈₈-Xaa₈₉-Xaa₉₀-Phe-Pro-Xaa₉₁-Leu-Glu-Xaa₉₂-Leu-
 Xaa₉₃-Φaa₂₁-Ωaa₁₅-Ωaa₁₆-Met-Pro-Σaa₁₆-Leu-Ωaa₁₇-Glu-Φaa₂₂

- wherein: each of Φ₁₋₂₂ is a hydrophobic amino acid residue,
 20 each of Σaa₁₋₁₆ is a small amino acid residue,
 each of Baa₁₋₁₄ is a basic amino acid residue,
 each of Åaa₁₋₂ is an acidic amino acid residue,
 each of Ωaa₁₋₁₆ is a charged amino acid residue, and
 Xaa₁₋₉₃ are each independently selected from any amino acid residue.
- 25 [0057] In some embodiments, Ωaa₁ is selected from His or Asp. In some
 embodiments, Ωaa₂ is selected from Asp or Arg. In some embodiments, Ωaa₃ is selected
 from Lys or Asp. In some embodiments, Ωaa₄ is selected from Glu or Lys. In some
 embodiments, Ωaa₅ is selected from Glu or Arg. In some embodiments, Ωaa₆ is selected
 from His or Asp. In some embodiments, Ωaa₇ is selected from Arg or Glu. In some
 30 embodiments, Ωaa₈ is selected from Glu or Lys. In some embodiments, Ωaa₉ is selected
 from Glu or Arg.

[0058] In some embodiments, Ω_{aa10} is selected from Glu or Arg. In some
embodiments, Ω_{aa11} is selected from Glu or Lys. In some embodiments, Ω_{aa12} is
selected from Lys or Asp. In some embodiments, Ω_{aa13} is selected from Lys or Glu. In
some embodiments, Ω_{aa14} is selected from Lys or Glu. In some embodiments, Ω_{aa15} is
5 selected from Glu or Arg. In some embodiments, Ω_{aa16} is selected from Asp or Arg.

[0059] In some embodiments, Φ_{aa1} is selected from Leu or Met. In some
embodiments, Φ_{aa2} is selected from Leu or Phe. In some embodiments, Φ_{aa3} is selected
from Met or Ile. In some embodiments, Φ_{aa4} is selected from Ile or Val. In some
embodiments, Φ_{aa5} is selected from Ile or Leu. In some embodiments, Φ_{aa6} is selected
10 from Leu or Val. In some embodiments, Φ_{aa7} is selected from Phe or Leu. In some
embodiments, Φ_{aa8} is selected from Ile or Leu. In some embodiments, Φ_{aa9} is selected
from Ile or Val.

[0060] In some embodiments, Φ_{aa10} is selected from Phe or Tyr. In some
embodiments, Φ_{aa11} is selected from Leu or Ile. In some embodiments, Φ_{aa12} is
15 selected from Leu or Met. In some embodiments, Φ_{aa13} is selected from Leu or Met. In
some embodiments, Φ_{aa14} is selected from Met or Tyr. In some embodiments, Φ_{aa15} is
selected from Leu or Met. In some embodiments, Φ_{aa16} is selected from Leu or Ile. In
some embodiments, Φ_{aa17} is selected from Ile or Leu. In some embodiments, Φ_{aa18} is
selected from Met or Ile. In some embodiments, Φ_{aa19} is selected from Met or Ile.

20 [0061] In some embodiments, Φ_{aa20} is selected from Leu or Phe. In some
embodiments, Φ_{aa21} is selected from Leu or Ile. In some embodiments, Φ_{aa22} is
selected from Phe or Trp.

[0062] In some embodiments, Baa_1 , Baa_{3-6} and Baa_{9-14} are each independently
selected from Arg or Lys. In some embodiments, Baa_2 and Baa_8 are each independently
25 selected from His or Arg. In some embodiments, Baa_2 is selected from His or Lys.

[0063] In some embodiments, Σ_{aa1} is selected from Ala or Thr. In some
embodiments, Σ_{aa2} is selected from Ala or Thr. In some embodiments, Σ_{aa3} is selected
from Gly or Ser. In some embodiments, Σ_{aa4} is selected from Ser or Ala. In some
embodiments, Σ_{aa5} is selected from Gly or Ala. In some embodiments, Σ_{aa6} is selected
30 from Gly or Ser. In some embodiments, Σ_{aa7} is selected from Ala or Ser. In some

embodiments, Σaa_8 is selected from Thr or Ser. In some embodiments, Σaa_9 is selected from Ser or Gly.

[0064] In some embodiments, Σaa_{10} is selected from Ala or Thr. In some embodiments, Σaa_{11} is selected from Thr or Ser. In some embodiments, Σaa_{12} is selected from Thr or Ala. In some embodiments, Σaa_{13} is selected from Ser or Pro. In some embodiments, Σaa_{14} is selected from Pro or Ser. In some embodiments, Σaa_{15} is selected from Thr or Gly. In some embodiments, Σaa_{16} is selected from Thr or Ser.

[0065] In some embodiments, each of Δaa_{1-2} is independently selected from Asp or Glu.

10 [0066] In some embodiments Xaa_1 is a small or hydrophobic amino acid residue, e.g., Xaa_1 is selected from Pro or Leu. In some embodiments, Xaa_2 is an small or basic amino acid residue, e.g., Xaa_2 is selected from Ser or Arg. In some embodiments, Xaa_3 is a basic or hydrophobic amino acid residue, e.g., Xaa_3 is selected from Arg or Met. In some embodiments, Xaa_4 is an basic or small amino acid residue, e.g., Xaa_4 is selected from Lys or Ser. In some embodiments, Xaa_5 is a hydrophobic or acidic amino acid residue, e.g., Xaa_5 is selected from Val or Asp. In some embodiments, Xaa_6 is a neutral/polar or small amino acid residue, e.g., Xaa_6 is selected from Gln or Ser. In some embodiments, Xaa_7 is a basic or neutral/polar amino acid residue, e.g., Xaa_7 is selected from Lys or Asn. In some embodiments, Xaa_8 is a small or hydrophobic amino acid residue, e.g., Xaa_8 is selected from Gly or Val. In some embodiments, Xaa_9 is an acidic or neutral/polar amino acid residue, e.g., Xaa_9 is selected from Glu or Asn.

[0067] In some embodiments, Xaa_{10} is an acidic or small amino acid residue, e.g., Xaa_{10} is selected from Asp or Ser. In some embodiments, Xaa_{11} is a hydrophobic or small amino acid residue, e.g., Xaa_{11} is selected from Ile or Ser. In some embodiments, Xaa_{12} is an acidic or neutral/polar amino acid residue, e.g., Xaa_{12} is selected from Asp or Asn. In some embodiments, Xaa_{13} is a hydrophobic or basic amino acid residue, e.g., Xaa_{13} is selected from Ile or Lys. In some embodiments, Xaa_{14} is a neutral/polar or basic amino acid residue, e.g., Xaa_{14} is selected from Gln or His. In some embodiments, Xaa_{15} is acidic or small amino acid residue, e.g., Xaa_{15} is selected from Asp or Gly. In some embodiments, Xaa_{16} is absent or is a hydrophobic amino acid residue, e.g., Tyr. In some embodiments, Xaa_{17} is a neutral/polar or basic amino acid residue, e.g., Xaa_{17} is selected from Cys or Arg. In some embodiments, Xaa_{18} is a neutral/polar or small

amino acid residue, e.g., Xaa₁₈ is selected from Cys or Thr. In some embodiments, Xaa₁₉ is a hydrophobic or neutral/polar amino acid residue, e.g., Xaa₁₉ is selected from Tyr or Cys.

[0068] In some embodiments, Xaa₂₀ is a neutral/polar or hydrophobic amino acid residue, e.g., Xaa₂₀ is selected from Asn or Leu. In some embodiments, Xaa₂₁ is a basic or hydrophobic amino acid residue, e.g., Xaa₂₁ is selected from Arg or Leu. In some embodiments, Xaa₂₂ is a hydrophobic or acidic amino acid residue, e.g., Xaa₂₂ is selected from Trp or Glu. In some embodiments, Xaa₂₃ is a neutral/polar or acidic amino acid residue, e.g., Xaa₂₃ is selected from Gln or Glu. In some embodiments, Xaa₂₄ is a basic or neutral/polar amino acid residue, e.g., Xaa₂₄ is selected from Arg or Cys. In some embodiments, Xaa₂₅ is a small or basic amino acid residue, e.g., Xaa₂₅ is selected from Ser or Arg. In some embodiments, Xaa₂₆ is a neutral/polar or basic amino acid residue, e.g., Xaa₂₆ is selected from Gln or Arg. In some embodiments, Xaa₂₇ is a neutral/polar or basic amino acid residue, e.g., Xaa₂₇ is selected from Asn or Lys. In some embodiments, Xaa₂₈ is a hydrophobic or small amino acid residue, e.g., Xaa₂₈ is selected from Val or Ala. In some embodiments, Xaa₂₉ is an acidic or neutral/polar amino acid residue, e.g., Xaa₂₉ is selected from Glu or Asn.

[0069] In some embodiments, Xaa₃₀ is an acidic or small amino acid residue, e.g., Xaa₃₀ is selected from Asp or Pro. In some embodiments, Xaa₃₁ is a hydrophobic or small amino acid residue, e.g., Xaa₃₁ is selected from Tyr or Ala. In some embodiments, Xaa₃₂ is a small or acidic amino acid residue, e.g., Xaa₃₂ is selected from Ser or Glu. In some embodiments, Xaa₃₃ is a small or basic amino acid residue, e.g., Xaa₃₃ is selected from Ser or Lys. In some embodiments, Xaa₃₄ is a basic or neutral/polar amino acid residue, e.g., Xaa₃₄ is selected from Lys or Asn. In some embodiments, Xaa₃₅ is a hydrophobic or acidic amino acid residue, e.g., Xaa₃₅ is selected from Leu or Asp. In some embodiments, Xaa₃₆ is a neutral/polar or basic amino acid residue, e.g., Xaa₃₆ is selected from Asn or His. In some embodiments, Xaa₃₇ is a neutral/polar or basic amino acid residue, e.g., Xaa₃₇ is selected from Asn or Lys. In some embodiments, Xaa₃₈ is a neutral/polar or basic amino acid residue, e.g., Xaa₃₈ is selected from Asn or Lys. In some embodiments, Xaa₃₉ is a basic or small amino acid residue, e.g., Xaa₃₉ is selected from Lys or Gly.

[0070] In some embodiments, Xaa₄₀ is a small or neutral/polar amino acid residue, e.g., Xaa₄₀ is selected from Thr or Asn. In some embodiments, Xaa₄₁ is a small or acidic

amino acid residue, e.g., Xaa₄₁ is selected from Thr or Asp. In some embodiments, Xaa₄₂ is a basic or small amino acid residue, e.g., Xaa₄₂ is selected from Arg or Ser. In some embodiments, Xaa₄₃ is a small or basic amino acid residue, e.g., Xaa₄₃ is selected from Thr or Arg. In some embodiments, Xaa₄₄ is an acidic or neutral/polar amino acid residue, e.g., Xaa₄₄ is selected from Glu or Gln. In some embodiments, Xaa₄₅ is a small or acidic amino acid residue, e.g., Xaa₄₅ is selected from Gly or Glu. In some embodiments, Xaa₄₆ is a small or basic amino acid residue, e.g., Xaa₄₆ is selected from Ser or Lys. In some embodiments, Xaa₄₇ is a basic or small amino acid residue, e.g., Xaa₄₇ is selected from Lys or Thr. In some embodiments, Xaa₄₈ is a small or basic amino acid residue, e.g., Xaa₄₈ is selected from Ser or Arg. In some embodiments, Xaa₄₉ is a hydrophobic or basic amino acid residue, e.g., Xaa₄₉ is selected from Tyr or Lys.

[0071] In some embodiments, Xaa₅₀ is a small or hydrophobic amino acid residue, e.g., Xaa₅₀ is selected from Ala or Leu. In some embodiments, X₅₁ is a hydrophobic or acidic amino acid residue, e.g., Xaa₅₁ is selected from Ala or Asp. In some embodiments, Xaa₅₂ is a neutral/polar or basic amino acid residue, e.g., Xaa₅₂ is selected from Gln or Arg. In some embodiments, Xaa₅₃ is a hydrophobic or small amino acid residue, e.g., Xaa₅₃ is selected from Val or Gly. In some embodiments, Xaa₅₄ is absent or is a small amino acid residue, e.g., Ser. In some embodiments, Xaa₅₅ is absent or is a small amino acid residue, e.g., Ser. In some embodiments, Xaa₅₆ is absent or is a hydrophobic amino acid residue, e.g., Leu. In some embodiments, Xaa₅₇ is an acidic or hydrophobic amino acid residue, e.g., Xaa₅₇ is selected from Glu or Ala. In some embodiments, Xaa₅₈ is a basic or small amino acid residue, e.g., Xaa₅₈ is selected from His or Gly. In some embodiments, Xaa₅₉ is a hydrophobic or neutral/polar amino acid residue, e.g., Xaa₅₉ is selected from Leu or Cys.

[0072] In some embodiments, Xaa₆₀ is a hydrophobic or acidic amino acid residue, e.g., Xaa₆₀ is selected from Leu or Asp. In some embodiments, X₆₁ is a hydrophobic or basic amino acid residue, e.g., Xaa₆₁ is selected from Val or Arg. In some embodiments, Xaa₆₂ is a small or acidic amino acid residue, e.g., Xaa₆₂ is selected from Ser or Asp. In some embodiments, Xaa₆₃ is a hydrophobic or basic amino acid residue, e.g., Xaa₆₃ is selected from Leu or Lys. In some embodiments, Xaa₆₄ is a neutral/polar or basic amino acid residue, e.g., Xaa₆₄ is selected from Gln or Arg. In some embodiments, Xaa₆₅ is a basic or small amino acid residue, e.g., Xaa₆₅ is selected from His or Pro. In some

embodiments, Xaa₆₆ is a hydrophobic or neutral/polar amino acid residue, e.g., Xaa₆₆ is selected from Phe or Asn. In some embodiments, Xaa₆₇ is a small or acidic amino acid residue, e.g., Xaa₆₇ is selected from Ser or Glu. In some embodiments, Xaa₆₈ is a small or hydrophobic amino acid residue, e.g., Xaa₆₈ is selected from Gly or Tyr. In some
5 embodiments, Xaa₆₉ is a hydrophobic or small amino acid residue, e.g., Xaa₆₉ is selected from Val or Ser.

[0073] In some embodiments, Xaa₇₀ is an acidic or small amino acid residue, e.g., Xaa₇₀ is selected from Asp or Thr. In some embodiments, Xaa₇₁ is a hydrophobic or acidic amino acid residue, e.g., Xaa₇₁ is selected from Val or Asp. In some embodiments,
10 Xaa₇₂ is a basic or neutral/polar amino acid residue, e.g., Xaa₇₂ is selected from Lys or Gln. In some embodiments, Xaa₇₃ is a small or acidic amino acid residue, e.g., Xaa₇₃ is selected from Gly or Glu. In some embodiments, Xaa₇₄ is a neutral/polar or small amino acid residue, e.g., Xaa₇₄ is selected from Asn or Ser. In some embodiments, Xaa₇₅ is an acidic or small amino acid residue, e.g., Xaa₇₅ is selected from Glu or Thr. In some
15 embodiments, Xaa₇₆ is a small or acidic amino acid residue, e.g., Xaa₇₆ is selected from Gly or Glu. In some embodiments, Xaa₇₇ is a hydrophobic or neutral/polar amino acid residue, e.g., Xaa₇₇ is selected from Tyr or Cys. In some embodiments, Xaa₇₈ is a hydrophobic or basic amino acid residue, e.g., Xaa₇₈ is selected from Phe or His. In some embodiments, Xaa₇₉ is a basic or hydrophobic amino acid residue, e.g., Xaa₇₉ is
20 selected from His or Ile.

[0074] In some embodiments, Xaa₈₀ is a hydrophobic or basic amino acid residue, e.g., Xaa₈₀ is selected from Val or His. In some embodiments, Xaa₈₁ is a small or hydrophobic amino acid residue, e.g., Xaa₈₁ is selected from Arg or Gly. In some
embodiments, Xaa₈₂ is a hydrophobic or neutral/polar amino acid residue, e.g., Xaa₈₂ is
25 selected from Val or Gln. In some embodiments, Xaa₈₃ is a small or neutral/polar amino acid residue, e.g., Xaa₈₃ is selected from Ser or Asn. In some embodiments, Xaa₈₄ is a basic or hydrophobic amino acid residue, e.g., Xaa₈₄ is selected from His or Leu. In some embodiments, Xaa₈₅ is a neutral/polar or acidic amino acid residue, e.g., Xaa₈₅ is selected from Gln or Glu. In some embodiments, Xaa₈₆ is a neutral/polar or
30 hydrophobic amino acid residue, e.g., Xaa₈₆ is selected from Cys or Tyr. In some embodiments, Xaa₈₇ is a neutral/polar or small amino acid residue, e.g., Xaa₈₇ is selected from Cys or Thr. In some embodiments, Xaa₈₈ is a small or hydrophobic amino

acid residue, e.g., Xaa₈₈ is selected from Ser or Val. In some embodiments, Xaa₈₉ is a basic or small amino acid residue, e.g., Xaa₈₉ is selected from Lys or Ser.

[0075] In some embodiments, Xaa₉₀ is a hydrophobic or small amino acid residue, e.g., Xaa₉₀ is selected from Leu or Gly. In some embodiments, Xaa₉₁ is a basic or
5 hydrophobic amino acid residue, e.g., Xaa₉₁ is selected from Arg or Leu. In some
embodiments, Xaa₉₂ is an acidic or hydrophobic amino acid residue, e.g., Xaa₉₂ is
selected from Glu or Leu. In some embodiments, Xaa₉₃ is a hydrophobic or
neutral/polar amino acid residue, e.g., Xaa₉₃ is selected from Val or Asn.

[0076] In yet another aspect, the invention provides antigen-binding molecules that
10 are specifically immuno-interactive with a polypeptide or portion as broadly described
above.

BRIEF DESCRIPTION OF THE DRAWINGS

[0077] Figure 1 is a schematic representation of the location and structure of the eight main classes of plant disease resistance proteins.

5 [0078] Figure 2 is a diagrammatic representation showing an alignment of the amino acid sequences set forth in SEQ ID NO: 2 and 4 using ClustalW multiple alignment and the PAM250 similarity matrix as disclosed for example by Dayhoff *et al.* (1978) A model of evolutionary change in proteins. Matrices for determining distance relationships *In* M. O. Dayhoff, (ed.), Atlas of protein sequence and structure, Vol. 5, pp. 345-358, National Biomedical Research Foundation, Washington DC; and by
10 Gonnet *et al.*, 1992, *Science* 256(5062): 144301445.

[0079] Figure 3 is a photographic representation showing the migration of amplification products on an agarose gel following an RT-PCR using primers specific for each banana NBS class using template RNA from *M. acuminata ssp. malaccensis* resistant (R) or susceptible (S) plants. Total RNA was extracted from leaf or root tissue
15 and treated with DNAase. C+ lanes, expected ~ 480 bp actin 1 cDNA fragment; C- lanes, no reverse transcriptase; AD, expected ~580 bp actin 1 genomic DNA fragment with ~100 bp intron included.

TABLE A

BRIEF DESCRIPTION OF THE SEQUENCES

SEQUENCE ID NUMBER	SEQUENCE	LENGTH
SEQ ID NO: 1	Nucleotide sequence of RGA5 obtained from Calcutta 4	4380 nts
SEQ ID NO: 2	Deduced amino acid sequence encoded by SEQ ID NO: 1	1441 aa
SEQ ID NO: 3	Nucleotide sequence of RGA2 obtained from <i>Musa acuminata</i> spp <i>malaccensis</i>	3699 nts
SEQ ID NO: 4	Deduced amino acid sequence encoded by SEQ ID NO: 1	1232 aa

DETAILED DESCRIPTION OF THE INVENTION

1. Definitions

[0080] Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by those of ordinary skill in the art to which
5 the invention belongs. Although any methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present invention, preferred methods and materials are described. For the purposes of the present invention, the following terms are defined below.

[0081] The articles "a" and "an" are used herein to refer to one or to more than one
10 (i.e., to at least one) of the grammatical object of the article. By way of example, "an element" means one element or more than one element.

[0082] By "about" is meant a quantity, level, value, frequency, percentage, dimension, size, amount, weight or length that varies by as much as 30, 25, 20, 25, 10,
15 9, 8, 7, 6, 5, 4, 3, 2 or 1 % to a reference quantity, level, value, frequency, percentage, dimension, size, amount, weight or length.

[0083] By "*antigen-binding molecule*" is meant a molecule that has binding affinity for a target antigen. It will be understood that this term extends to immunoglobulins, immunoglobulin fragments and non-immunoglobulin derived protein frameworks that exhibit antigen-binding activity.

20 [0084] As used herein, the term "*binds specifically*," "*specifically immuno-interactive*" and the like refers to antigen-binding molecules that bind or a immuno-interactive with the polypeptide or polypeptide portions of the invention but do not significantly bind to homologous prior art polypeptides.

[0085] By "*biologically active portion*" is meant a portion of a full-length parent
25 peptide or polypeptide which portion retains an activity of the parent molecule. For example, a biologically active portion of polypeptide of the invention will retain the ability to confer disease resistance, especially resistance to fungal pathogens such as *Fusarium*. As used herein, the term "*biologically active portion*" includes deletion mutants and peptides, for example of at least about 8, 9, 10, 11, 12, 13, 14, 15, 16, 17,
30 18, 19, 20, 21, 22, 23, 24, 25, 30, 40, 50, 60, 70, 80, 90, 100, 120, 150, 300, 400, 500, 600, 700, 800, 900 or 1000 contiguous amino acids, which comprise an activity of a parent molecule. Portions of this type may be obtained through the application of

standard recombinant nucleic acid techniques or synthesised using conventional liquid or solid phase synthesis techniques. For example, reference may be made to solution synthesis or solid phase synthesis as described, for example, in Chapter 9 entitled "Peptide Synthesis" by Atherton and Shephard which is included in a publication 5 entitled "Synthetic Vaccines" edited by Nicholson and published by Blackwell Scientific Publications. Alternatively, peptides can be produced by digestion of a peptide or polypeptide of the invention with proteinases such as endoLys-C, endoArg-C, endoGlu-C and staphylococcus V8-protease. The digested fragments can be purified by, for example, high performance liquid chromatographic (HPLC) techniques. 10 Recombinant nucleic acid techniques can also be used to produce such portions.

[0086] As used herein, the term "*cis-acting sequence*", "*cis-acting element*" or "*cis-regulatory region*" or "*regulatory region*" or similar term shall be taken to mean any sequence of nucleotides, which when positioned appropriately relative to an expressible genetic sequence, is capable of regulating, at least in part, the expression of the genetic 15 sequence. Those skilled in the art will be aware that a *cis-regulatory region* may be capable of activating, silencing, enhancing, repressing or otherwise altering the level of expression and/or cell-type-specificity and/or developmental specificity of a gene sequence at the transcriptional or post-transcriptional level. In certain embodiments of the present invention, the *cis-acting sequence* is an activator sequence that enhances or 20 stimulates the expression of an expressible genetic sequence. Throughout this specification, unless the context requires otherwise, the words "*comprise*", "*comprises*" and "*comprising*" will be understood to imply the inclusion of a stated step or element or group of steps or elements but not the exclusion of any other step or element or group of steps or elements.

25 [0087] By "*corresponds to*" or "*corresponding to*" is meant a polynucleotide (a) having a nucleotide sequence that is substantially identical or complementary to all or a portion of a reference polynucleotide sequence or (b) encoding an amino acid sequence identical to an amino acid sequence in a peptide or protein. This phrase also includes within its scope a peptide or polypeptide having an amino acid sequence that is 30 substantially identical to a sequence of amino acids in a reference peptide or protein.

[0088] As used herein, the terms "*culturing*", "*culture*" and the like refer to the set of procedures used *in vitro* where a population of cells (or a single cell) is incubated under conditions which have been shown to support the growth or maintenance of the

cells *in vitro*. The art recognises a wide number of formats, media, temperature ranges, gas concentrations etc. which need to be defined in a culture system. The parameters will vary based on the format selected and the specific needs of the individual who practices the methods herein disclosed. However, it is recognised that the determination
5 of culture parameters is routine in nature.

[0089] By "*disease resistance*" is intended that plants avoid or suppress the disease symptoms that are the outcome of plant-pathogen interaction. That is, pathogens are prevented from causing plant diseases and the associated disease symptoms. The methods of the invention can be utilised to protect plants from disease, particularly
10 those diseases that are caused by plant pathogens, such as Fusarium wilt.

[0090] By "*expression vector*" is meant any autonomous genetic element capable of directing the transcription of a polynucleotide contained within the vector and suitably the synthesis of a peptide or polypeptide encoded by the polynucleotide. Such expression vectors are known to practitioners in the art.

15 [0091] The term "*gene*" as used herein refers to any and all discrete coding regions of the cell's genome, as well as associated non-coding and regulatory regions. The gene is also intended to mean the open reading frame encoding specific polypeptides, introns, and adjacent 5' and 3' non-coding nucleotide sequences involved in the regulation of expression. In this regard, the gene may further comprise control signals such as
20 promoters, enhancers, termination and/or polyadenylation signals that are naturally associated with a given gene, or heterologous control signals. The DNA sequences may be cDNA or genomic DNA or a fragment thereof. The gene may be introduced into an appropriate vector for extrachromosomal maintenance or for integration into the host.

[0092] The terms "*growing*" or "*regeneration*" as used herein mean growing a
25 whole, differentiated plant from a plant cell, a group of plant cells, a plant part (including seeds), or a plant piece (e.g., from a protoplast, callus, or tissue part).

[0093] "*Hybridisation*" is used herein to denote the pairing of complementary nucleotide sequences to produce a DNA-DNA hybrid or a DNA-RNA hybrid. Complementary base sequences are those sequences that are related by the base-pairing
30 rules. In DNA, A pairs with T and C pairs with G. In RNA U pairs with A and C pairs with G. In this regard, the terms "match" and "mismatch" as used herein refer to the hybridisation potential of paired nucleotides in complementary nucleic acid strands. Matched nucleotides hybridise efficiently, such as the classical A-T and G-C base pair

mentioned above. Mismatches are other combinations of nucleotides that do not hybridise efficiently.

[0094] Reference herein to "*immuno-interactive*" includes reference to any interaction, reaction, or other form of association between molecules and in particular

5 where one of the molecules is, or mimics, a component of the immune system.

[0095] By "*isolated*" is meant material that is substantially or essentially free from components that normally accompany it in its native state. For example, an "*isolated polynucleotide*", as used herein, refers to a polynucleotide, which has been purified from the sequences which flank it in a naturally-occurring state, e.g., a DNA fragment

10 which has been removed from the sequences that are normally adjacent to the fragment. Alternatively, an "*isolated peptide*" or an "*isolated polypeptide*" and the like, as used herein, refer to *in vitro* isolation and/or purification of a peptide or polypeptide molecule from its natural cellular environment, and from association with other components of the cell, i.e., it is not associated with *in vivo* substances.

15 [0096] By "*marker gene*" is meant a gene that imparts a distinct phenotype to cells expressing the marker gene and thus allows such transformed cells to be distinguished from cells that do not have the marker. A selectable marker gene confers a trait for which one can 'select' based on resistance to a selective agent (e.g., a herbicide, antibiotic, radiation, heat, or other treatment damaging to untransformed cells). A
20 screenable marker gene (or reporter gene) confers a trait that one can identify through observation or testing, i.e., by 'screening' (e.g. β -glucuronidase, luciferase, or other enzyme activity not present in untransformed cells).

[0097] As used herein, a "*naturally-occurring*" nucleic acid molecule refers to an RNA or DNA molecule having a nucleotide sequence that occurs in nature. For
25 example a naturally-occurring nucleic acid molecule can encode a natural protein.

[0098] By "*obtained from*." is meant that a sample such as, for example, a nucleic acid extract or polypeptide extract is isolated from, or derived from, a particular source. For example, the extract may be isolated directly from any membrane-translocating sequence-containing organism, such as but not limited to bacteria, yeast and plants as
30 well as animals including mammals, birds, reptiles, fish and insects.

[0099] The term "*oligonucleotide*" as used herein refers to a polymer composed of a multiplicity of nucleotide residues (deoxyribonucleotides or ribonucleotides, or related

structural variants or synthetic analogues thereof, including nucleotides with modified or substituted sugar groups and the like) linked *via* phosphodiester bonds (or related structural variants or synthetic analogues thereof). Thus, while the term "oligonucleotide" typically refers to a nucleotide polymer in which the nucleotide residues and linkages between them are naturally-occurring, it will be understood that the term also includes within its scope various analogues including, but not restricted to, peptide nucleic acids (PNAs), phosphorothioate, phosphorodithioate, phosphoroselenoate, phosphorodiselenoate, phosphoroanilothioate, phosphoraniladate, phosphoroamidate, methyl phosphonates, 2-O-methyl ribonucleic acids, and the like.

The exact size of the molecule can vary depending on the particular application. Oligonucleotides are a polynucleotide subset with 200 bases or fewer in length. Preferably, oligonucleotides are 10 to 60 bases in length and most preferably 12, 13, 14, 15, 16, 17, 18, 19, or 20 to 40 bases in length. Oligonucleotides are usually single stranded, e.g., for probes; although oligonucleotides may be double stranded, e.g., for use in the construction of a variant nucleic acid sequence. Oligonucleotides of the invention can be either sense or antisense oligonucleotides.

[00100] The term "*operably connected*" or "*operably linked*" as used herein means placing a structural gene under the regulatory control of a promoter, which then controls the transcription and optionally translation of the gene. In the construction of heterologous promoter/structural gene combinations, it is generally preferred to position the genetic sequence or promoter at a distance from the gene transcription start site that is approximately the same as the distance between that genetic sequence or promoter and the gene it controls in its natural setting; i.e. the gene from which the genetic sequence or promoter is derived. As is known in the art, some variation in this distance can be accommodated without loss of function. Similarly, the preferred positioning of a regulatory sequence element with respect to a heterologous gene to be placed under its control is defined by the positioning of the element in its natural setting; i.e., the genes from which it is derived.

[0100] The term "*pathogen*" is used herein in its broadest sense to refer to an organism or an infectious agent whose infection of cells of viable plant tissue elicits a disease response.

[0101] The term "*polynucleotide*" or "*nucleic acid*" as used herein designates mRNA, RNA, cRNA, cDNA or DNA. The term typically refers polymeric form of

nucleotides of at least 10 bases in length, either ribonucleotides or deoxynucleotides or a modified form of either type of nucleotide. The term includes single and double stranded forms of DNA.

5 [0102] The terms "*polynucleotide variant*" and "*variant*" refer to polynucleotides displaying substantial sequence identity with a reference polynucleotide sequence or polynucleotides that hybridise with a reference sequence under stringent conditions that are defined hereinafter. These terms also encompass polynucleotides in which one or more nucleotides have been added or deleted, or replaced with different nucleotides. In this regard, it is well understood in the art that certain alterations inclusive of mutations, 10 additions, deletions and substitutions can be made to a reference polynucleotide whereby the altered polynucleotide retains a biological function or activity of the reference polynucleotide. The terms "*polynucleotide variant*" and "*variant*" also include naturally-occurring allelic variants.

[0103] "*Polypeptide*", "*peptide*" and "*protein*" are used interchangeably herein to 15 refer to a polymer of amino acid residues and to variants and synthetic analogues of the same. Thus, these terms apply to amino acid polymers in which one or more amino acid residues is a synthetic non-naturally-occurring amino acid, such as a chemical analogue of a corresponding naturally-occurring amino acid, as well as to naturally-occurring amino acid polymers.

20 [0104] The term "*polypeptide variant*" refers to polypeptides which are distinguished from a reference polypeptide by the addition, deletion or substitution of at least one amino acid residue. In certain embodiments, one or more amino acid residues of a reference polypeptide are replaced by different amino acids. It is well understood in the art that some amino acids may be changed to others with broadly similar properties 25 without changing the nature of the activity of the polypeptide (conservative substitutions) as described hereinafter.

[0105] By "*primer*" is meant an oligonucleotide which, when paired with a strand of DNA, is capable of initiating the synthesis of a primer extension product in the presence of a suitable polymerising agent. The primer is preferably single-stranded for 30 maximum efficiency in amplification but can alternatively be double-stranded. A primer must be sufficiently long to prime the synthesis of extension products in the presence of the polymerisation agent. The length of the primer depends on many factors, including application, temperature to be employed, template reaction conditions, other reagents,

and source of primers. For example, depending on the complexity of the target sequence, the oligonucleotide primer typically contains 15 to 35 or more nucleotide residues, although it can contain fewer nucleotide residues. Primers can be large polynucleotides, such as from about 35 nucleotides to several kilobases or more.

- 5 Primers can be selected to be "substantially complementary" to the sequence on the template to which it is designed to hybridise and serve as a site for the initiation of synthesis. By "substantially complementary", it is meant that the primer is sufficiently complementary to hybridise with a target polynucleotide. Preferably, the primer contains no mismatches with the template to which it is designed to hybridise but this is
- 10 not essential. For example, non-complementary nucleotide residues can be attached to the 5' end of the primer, with the remainder of the primer sequence being complementary to the template. Alternatively, non-complementary nucleotide residues or a stretch of non-complementary nucleotide residues can be interspersed into a primer, provided that the primer sequence has sufficient complementarity with the sequence of
- 15 the template to hybridise therewith and thereby form a template for synthesis of the extension product of the primer.

- [0106] "Probe" refers to a molecule that binds to a specific sequence or sub-sequence or other moiety of another molecule. Unless otherwise indicated, the term "probe" typically refers to a polynucleotide probe that binds to another polynucleotide,
- 20 often called the "target polynucleotide", through complementary base pairing. Probes can bind target polynucleotides lacking complete sequence complementarity with the probe, depending on the stringency of the hybridisation conditions. Probes can be labelled directly or indirectly.

- [0107] The term "*recombinant polynucleotide*" as used herein refers to a
- 25 polynucleotide formed *in vitro* by the manipulation of nucleic acid into a form not normally found in nature. For example, the recombinant polynucleotide may be in the form of an expression vector. Generally, such expression vectors include transcriptional and translational regulatory nucleic acid operably linked to the nucleotide sequence.

- [0108] By "*recombinant polypeptide*" is meant a polypeptide made using
- 30 recombinant techniques, i.e., through the expression of a recombinant or synthetic polynucleotide.

- [0109] By "*regulatory element*" or "*regulatory element*" is meant nucleic acid sequences (e.g., DNA) necessary for expression of an operably linked coding sequence

in a particular host cell. The regulatory sequences that are suitable for prokaryotic cells for example, include a promoter, and optionally a *cis*-acting sequence such as an operator sequence and a ribosome binding site. Control sequences that are suitable for eukaryotic cells include promoters, polyadenylation signals, transcriptional enhancers, translational enhancers, leader or trailing sequences that modulate mRNA stability, as well as targeting sequences that target a product encoded by a transcribed polynucleotide to an intracellular compartment within a cell or to the extracellular environment.

[0110] The term "*sequence identity*" as used herein refers to the extent that sequences are identical on a nucleotide-by-nucleotide basis or an amino acid-by-amino acid basis over a window of comparison. Thus, a "*percentage of sequence identity*" is calculated by comparing two optimally aligned sequences over the window of comparison, determining the number of positions at which the identical nucleic acid base (e.g., A, T, C, G, I) or the identical amino acid residue (e.g., Ala, Pro, Ser, Thr, Gly, Val, Leu, Ile, Phe, Tyr, Trp, Lys, Arg, His, Asp, Glu, Asn, Gln, Cys and Met) occurs in both sequences to yield the number of matched positions, dividing the number of matched positions by the total number of positions in the window of comparison (i.e., the window size), and multiplying the result by 100 to yield the percentage of sequence identity. For the purposes of the present invention, "*sequence identity*" will be understood to mean the "match percentage" calculated by the DNASIS computer program (Version 2.5 for windows; available from Hitachi Software engineering Co., Ltd., South San Francisco, California, USA) using standard defaults as used in the reference manual accompanying the software.

[0111] "*Similarity*" refers to the percentage number of amino acids that are identical or constitute conservative substitutions as defined in Table B *infra*. Similarity may be determined using sequence comparison programs such as GAP (Deveraux *et al.* 1984, *Nucleic Acids Research* 12, 387-395). In this way, sequences of a similar or substantially different length to those cited herein might be compared by insertion of gaps into the alignment, such gaps being determined, for example, by the comparison algorithm used by GAP.

[0112] Terms used to describe sequence relationships between two or more polynucleotides or polypeptides include "reference sequence", "comparison window", "sequence identity", "percentage of sequence identity" and "substantial identity". A

"reference sequence" is at least 12 but frequently 15 to 18 and often at least 25 monomer units, inclusive of nucleotides and amino acid residues, in length. Because two polynucleotides may each comprise (1) a sequence (i.e., only a portion of the complete polynucleotide sequence) that is similar between the two polynucleotides, and
5 (2) a sequence that is divergent between the two polynucleotides, sequence comparisons between two (or more) polynucleotides are typically performed by comparing sequences of the two polynucleotides over a "comparison window" to identify and compare local regions of sequence similarity. A "comparison window" refers to a conceptual segment of at least 6 contiguous positions, usually about 50 to about 100,
10 more usually about 100 to about 150 in which a sequence is compared to a reference sequence of the same number of contiguous positions after the two sequences are optimally aligned. The comparison window may comprise additions or deletions (i.e., gaps) of about 20% or less as compared to the reference sequence (which does not comprise additions or deletions) for optimal alignment of the two sequences. Optimal
15 alignment of sequences for aligning a comparison window may be conducted by computerised implementations of algorithms (GAP, BESTFIT, FASTA, and TFASTA in the Wisconsin Genetics Software Package Release 7.0, Genetics Computer Group, 575 Science Drive Madison, WI, USA) or by inspection and the best alignment (i.e., resulting in the highest percentage homology over the comparison window) generated
20 by any of the various methods selected. Reference also may be made to the BLAST family of programs as for example disclosed by Altschul *et al.*, 1997, *Nucl. Acids Res.* 25:3389. A detailed discussion of sequence analysis can be found in Unit 19.3 of Ausubel *et al.*, "Current Protocols in Molecular Biology", John Wiley & Sons Inc, 1994-1998, Chapter 15.

25 [0113] By the term "*taxon*" herein is meant a unit of botanical classification. It thus includes, genus, species, cultivars, varieties, variants and other minor taxonomic groups which lack a consistent nomenclature.

[0114] The term "*transformation*" means alteration of the genotype of an organism, for example a bacterium, yeast or plant, by the introduction of a foreign or endogenous
30 nucleic acid.

[0115] By "*vector*" is meant a polynucleotide molecule, preferably a DNA molecule derived, for example, from a plasmid, bacteriophage, yeast or virus, into which a polynucleotide can be inserted or cloned. A vector preferably contains one or

more unique restriction sites and can be capable of autonomous replication in a defined host cell including a target cell or tissue or a progenitor cell or tissue thereof, or be integrable with the genome of the defined host such that the cloned sequence is reproducible. Accordingly, the vector can be an autonomously replicating vector, i.e., a vector that exists as an extrachromosomal entity, the replication of which is independent of chromosomal replication, e.g., a linear or closed circular plasmid, an extrachromosomal element, a minichromosome, or an artificial chromosome. The vector can contain any means for assuring self-replication. Alternatively, the vector can be one which, when introduced into the host cell, is integrated into the genome and replicated together with the chromosome(s) into which it has been integrated. A vector system can comprise a single vector or plasmid, two or more vectors or plasmids, which together contain the total DNA to be introduced into the genome of the host cell, or a transposon. The choice of the vector will typically depend on the compatibility of the vector with the host cell into which the vector is to be introduced. In the present case, the vector is preferably a viral or viral-derived vector, which is operably functional in animal and preferably mammalian cells. Such vector may be derived from a poxvirus, an adenovirus or yeast. The vector can also include a selection marker such as an antibiotic resistance gene that can be used for selection of suitable transformants. Examples of such resistance genes are known to those of skill in the art.

2. Modulation of disease resistance

[0116] The invention is drawn to polynucleotide, polypeptide and methods for modulating disease resistance, especially for stimulating or enhancing disease resistance in plants, caused by pathogens. Pathogens of the invention include, but are not limited to, viruses or viroids, bacteria, insects, nematodes, fungi, and the like. Viruses include any plant virus, for example, tobacco or cucumber mosaic virus, ringspot virus, necrosis virus, maize dwarf mosaic virus, etc. Specific fungal and viral pathogens for the major crops include:

[0117] Soybeans: *Phytophthora megasperma* fsp. *glycinea*, *Macrophomina phaseolina*, *Rhizoctonia solani*, *Sclerotinia sclerotiorum*, *Fusarium oxysporum*, *Diaporthe phaseolorum* var. *sojae* (*Phomopsis sojae*), *Diaporthe phaseolorum* var. *caulivora*, *Sclerotium rolfsii*, *Cercospora kikuchii*, *Cercospora sojae*, *Peronospora manshurica*, *Colletotrichum dematium* (*Colletotrichum truncatum*), *Corynespora cassicola*, *Septoria glycines*, *Phyllosticta sojicola*, *Alternaria alternata*, *Pseudomonas*

syringae p.v. *glycinea*, *Xanthomonas campestris* p.v. *phaseoli*, *Microsphaera diffusa*, *Fusarium semitectum*, *Phialophora gregata*, Soybean mosaic virus, *Glomerella glycines*, Tobacco Ring spot virus, Tobacco Streak virus, *Phakopsorapachyrhizi*, *Pythium aphanidermatum*, *Pythium ultimum*, *Pythium debaryanum*, Tomato spotted wilt virus, *Heterodera glycines* *Fusarium solani*;

[0118] Canola: *Albugo candida*, *Alternaria brassicae*, *Leptosphaeria maculans*, *Rhizoctonia solani*, *Sclerotinia sclerotiorum*, *Mycosphaerella brassicicola*, *Pythium ultimum*, *Peronospora parasitica*, *Fusarium roseum*, *Alternaria alternata*;

[0119] Alfalfa: *Clavibater michiganese* subsp. *insidiosum*, *Pythium ultimum*, *Pythium irregulare*, *Pythium splendens*, *Pythium debaryanum*, *Pythium aphanidermatum*, *Phytophthora megasperma*, *Peronospora trifoliorum*, *Phoma medicaginis* var. *medicaginis*, *Cercospora medicaginis*, *Pseudopeziza medicaginis*, *Leptotrochila medicaginis*, *Fusarium*, *Xanthomonas campestris* p.v. *alfalfae*, *Aphanomyces euteiches*, *Stemphylium herbarum*, *Stemphylium alfalfae*;

[0120] Wheat: *Pseudomonas syringae* p.v. *atrofaciens*, *Urocystis agropyri*, *Xanthomonas campestris* p.v. *translucens*, *Pseudomonas syringae* p.v. *syringae*, *Alternaria alternata*, *Cladosporium herbarum*, *Fusarium graminearum*, *Fusarium avenaceum*, *Fusarium culmorum*, *Ustilago tritici*, *Ascochyta tritici*, *Cephalosporium gramineum*, *Collotetrichum graminicola*, *Erysiphe graminis* f.sp. *tritici*, *Puccinia graminis* f.sp. *tritici*, *Puccinia recondita* f.sp. *tritici*, *Puccinia striiformis*, *Pyrenophora tritici-repentis*, *Septoria nodorum*, *Septoria tritici*, *Septoria avenae*, *Pseudocercospora herpotrichoides*, *Rhizoctonia solani*, *Rhizoctonia cerealis*, *Gaeumannomyces graminis* var. *tritici*, *Pythium aphanidermatum*, *Pythium arrhenomanes*, *Pythium ultimum*, *Bipolaris sorokiniana*, Barley Yellow Dwarf Virus, Brome Mosaic Virus, Soil Borne Wheat Mosaic Virus, Wheat Streak Mosaic Virus, Wheat Spindle Streak Virus, American Wheat Striate Virus, *Claviceps purpurea*, *Tilletia tritici*, *Tilletia laevis*, *Ustilago tritici*, *Tilletia indica*, *Rhizoctonia solani*, *Pythium arrhenomanes*, *Pythium graminicola*, *Pythium aphanidermatum*, High Plains Virus, European wheat striate virus;

[0121] Sunflower: *Plasmophora halstedii*, *Sclerotinia sclerotiorum*, Aster Yellows, *Septoria helianthi*, *Phomopsis helianthi*, *Alternaria helianthi*, *Alternaria zinniae*, *Botrytis cinerea*, *Phoma macdonaldii*, *Macrophomina phaseolina*, *Erysiphe cichoracearum*, *Rhizopus oryzae*, *Rhizopus arrhizus*, *Rhizopus stolonifer*, *Puccinia*

helianthi, *Verticillium dahliae*, *Erwinia carotovorum* pv. *carotovora*, *Cephalosporium acremonium*, *Phytophthora cryptogea*, *Albugo tragopogonis*;

- [0122] Corn: *Fusarium moniliforme* var. *subglutinans*, *Erwinia stewartii*, *Fusarium moniliforme*, *Gibberella zeae* (*Fusarium graminearum*), *Stenocarpella maydi* (*Diplodia maydis*), *Pythium irregulare*, *Pythium debaryanum*, *Pythium graminicola*, *Pythium splendens*, *Pythium ultimum*, *Pythium aphanidermatum*, *Aspergillus flavus*, *Bipolaris maydis* O, T (*Cochliobolus heterostrophus*), *Helminthosporium carbonum* I, II & III (*Cochliobolus carbonum*), *Exserohilum turcicum* I, II & III, *Helminthosporium pedicellatum*, *Physoderma maydis*, *Phyllosticta maydis*, *Kabatiella-maydis*, *Cercospora sorghi*, *Ustilago maydis*, *Puccinia sorghi*, *Puccinia polysora*, *Macrophomina phaseolina*, *Penicillium oxalicum*, *Nigrospora oryzae*, *Cladosporium herbarum*, *Curvularia lunata*, *Curvularia inaequalis*, *Curvularia pallescens*, *Clavibacter michiganense* subsp. *nebraskense*, *Trichoderma viride*, Maize Dwarf Mosaic Virus A & B, Wheat Streak Mosaic Virus, Maize Chlorotic Dwarf Virus, *Claviceps sorghi*, *Pseudomonas avenae*, *Erwinia chrysanthemi* pv. *zea*, *Erwinia carotovora*, Corn stunt spiroplasma, *Diplodia macrospora*, *Sclerophthora macrospora*, *Peronosclerospora sorghi*, *Peronosclerospora philippinensis*, *Peronosclerospora maydis*, *Peronosclerospora sacchari*, *Sphacelotheca reiliana*, *Physopella zeae*, *Cephalosporium maydis*, *Cephalosporium acremonium*, Maize Chlorotic Mottle Virus, High Plains Virus, Maize Mosaic Virus, Maize Rayado Fino Virus, Maize Streak Virus, Maize Stripe Virus, Maize Rough Dwarf Virus;

- [0123] Sorghum: *Exserohilum turcicum*, *Colletotrichum graminicola* (*Glomerella graminicola*), *Cercospora sorghi*, *Gloeocercospora sorghi*, *Ascochyta sorghina*, *Pseudomonas syringae* p.v. *syringae*, *Xanthomonas campestris* p.v. *holcicola*, *Pseudomonas andropogonis*, *Puccinia purpurea*, *Macrophomina phaseolina*, *Perconia circinata*, *Fusarium moniliforme*, *Alternaria alternata*, *Bipolaris sorghicola*, *Helminthosporium sorghicola*, *Curvularia lunata*, *Phoma insidiosa*, *Pseudomonas avenae* (*Pseudomonas alboprecipitans*), *Ramulispora sorghi*, *Ramulispora sorghicola*, *Phyllachara sacchari*, *Sporisorium reilianum* (*Sphacelotheca reiliana*), *Sphacelotheca cruenta*, *Sporisorium sorghi*, Sugarcane mosaic H, Maize Dwarf Mosaic Virus A & B, *Claviceps sorghi*, *Rhizoctonia solani*, *Acremonium strictum*, *Sclerophthona macrospora*, *Peronosclerospora sorghi*, *Peronosclerospora philippinensis*, *Sclerospora*

graminicola, *Fusarium graminearum*, *Fusarium oxysporum*, *Pythium arrhenomanes*, *Pythium graminicola*, etc.

[0124] Nematodes include parasitic nematodes such as root-knot, cyst, lesion, and reniform nematodes, etc

5 [0125] Insect pests include insects selected from the orders Coleoptera, Diptera, Hymenoptera, Lepidoptera, Mallophaga, Homoptera, Hemiptera, Orthoptera, Thysanoptera, Dermaptera, Isoptera, Anoplura, Siphonaptera, Trichoptera, etc., particularly Coleoptera and Lepidoptera. Insect pests of the invention for the major crops include:

10 [0126] Maize: *Ostrinia nubilalis*, European corn borer; *Agrotis ipsilon*, black cutworm; *Helicoverpa zea*, corn earworm; *Spodoptera frugiperda*, fall armyworm; *Diatraea grandiosella*, southwestern corn borer; *Elasmopalpus lignosellus*, lesser cornstalk borer; *Diatraea saccharalis*, sugarcane borer; *Diabrotica virgifera*, western corn rootworm; *Diabrotica longicornis* barberi, northern corn rootworm; *Diabrotica*
 15 *undecimpunctata howardi*, southern corn rootworm; *Melanotus* spp., wireworms; *Cyclocephala borealis*, northern masked chafer (white grub); *Cyclocephala immaculata*, southern masked chafer (white grub); *Popillia japonica*, Japanese beetle; *Chaetocnema pulicaria*, corn flea beetle; *Sphenophorus maidis*, maize billbug; *Rhopalosiphum maidis*, corn leaf aphid; *Anuraphis maidiradicis*, corn root aphid;
 20 *Blissus leucopterus leucopterus*, chinch bug; *Melanoplus femurrubrum*, redlegged grasshopper; *Melanoplus sanguinipes*, migratory grasshopper; *Hylemya platura*, seedcorn maggot; *Agromyza parvicornis*, corn blot leafminer; *Anaphothrips obscurus*, grass thrips; *Solenopsis milesta*, thief ant; *Tetranychus urticae*, twospotted spider mite;

[0127] Sorghum: *Chilo partellus*, sorghum borer; *Spodoptera frugiperda*, fall
 25 armyworm; *Helicoverpa zea*, corn earworm; *Elasmopalpus lignosellus*, lesser cornstalk borer; *Feltia subterranea*, granulate cutworm; *Phyllophaga crinita*, white grub; *Eleodes*, *Conoderus*, and *Aeolus* spp., wireworms; *Oulema melanopus*, cereal leaf beetle; *Chaetocnema pulicaria*, corn flea beetle; *Sphenophorus maidis*, maize billbug; *Rhopalosiphum maidis*; corn leaf aphid; *Sipha flava*, yellow sugarcane aphid; *Blissus*
 30 *leucopterus leucopterus*, chinch bug; *Contarinia sorghicola*, sorghum midge; *Tetranychus cinnabarinus*, carmine spider mite; *Tetranychus urticae*, twospotted spider mite; Wheat: *Pseudaletia unipunctata*, army worm; *Spodoptera frugiperda*, fall armyworm; *Elasmopalpus lignosellus*, lesser cornstalk borer; *Agrotis orthogonia*,

- western cutworm; *Elasmopalpus lignosellus*, lesser cornstalk borer; *Oulema melanopus*, cereal leaf beetle; *Hypera punctata*, clover leaf weevil; *Diabrotica undecimpunctata* howardi, southern corn rootworm; Russian wheat aphid; *Schizaphis graminum*, greenbug; *Macrosiphum avenae*, English grain aphid; *Melanoplus femurrubrum*,
 5 redlegged grasshopper; *Melanoplus differentialis*, differential grasshopper; *Melanoplus sanguinipes*, migratory grasshopper; *Mayetiola destructor*, Hessian fly; *Sitodiplosis mosellana*, wheat midge; *Meromyza americana*, wheat stem maggot; *Hylemya coarctata*, wheat bulb fly; *Frankliniella fusca*, tobacco thrips; *Cephus cinctus*, wheat stem sawfly; *Aceria tulipae*, wheat curl mite; Sunflower: *Suleima helianthana*,
 10 sunflower bud moth; *Homoeosoma electellum*, sunflower moth; *zygogramma exclamationis*, sunflower beetle; *Bothyrus gibbosus*, carrot beetle; *Neolasioptera murtfeldtiana*, sunflower seed midge; Cotton: *Heliothis virescens*, cotton budworm; *Helicoverpa zea*, cotton bollworm; *Spodoptera exigua*, beet armyworm; *Pectinophora gossypiella*, pink bollworm; *Anthonomus grandis grandis*, boll weevil; *Aphis gossypii*,
 15 cotton aphid; *Pseudatomoscelis seriatus*, cotton fleahopper; *Trialeurodes abutilonea*, bandedwinged whitefly; *Lygus lineolaris*, tarnished plant bug; *Melanoplus femurrubrum*, redlegged grasshopper; *Melanoplus differentialis*, differential grasshopper; Thrips tabaci, onion thrips; *Frankliniella fusca*, tobacco thrips; *Tetranychus cinnabarinus*, carmine spider mite; *Tetranychus urticae*, twospotted spider mite;
- 20 [0128] Rice: *Diatraea saccharalis*, sugarcane borer; *Spodoptera frugiperda*, fall armyworm; *Helicoverpa zea*, corn earworm; *Colaspis brunnea*, grape colaspis; *Lissorhoptrus oryzophilus*, rice water weevil; *Sitophilus oryzae*, rice weevil; *Nephotettix nigropictus*, rice leafhopper; *Blissus leucopterus leucopterus*, chinch bug; *Acrosternum hilare*, green stink bug;
- 25 [0129] Soybean: *Pseudoplusia includens*, soybean looper; *Anticarsia gemmatilis*, velvetbean caterpillar; *Plathypena scabra*, green cloverworm; *Ostrinia nubilalis*, European corn borer; *Agrotis ipsilon*, black cutworm; *Spodoptera exigua*, beet armyworm; *Heliothis virescens*, cotton budworm; *Helicoverpa zea*, cotton bollworm; *Epilachna varivestis*, Mexican bean beetle; *Myzus persicae*, green peach aphid;
- 30 *Empoasca fabae*, potato leafhopper; *Acrosternum hilare*, green stink bug; *Melanoplus femurrubrum*, redlegged grasshopper; *Melanoplus differentialis*, differential grasshopper; *Hylemya platura*, seedcorn maggot; *Sericothrips variabilis*, soybean

thrips; *Thrips tabaci*, onion thrips; *Tetranychus turkestanii*, strawberry spider mite; *Tetranychus urticae*, twospotted spider mite;

[0130] Barley: *Ostrinia nubilalis*, European corn borer; *Agrotis ipsilon*, black cutworm; *Schizaphis graminum*, greenbug; *Blissus leucopterus leucopterus*, chinch bug; 5 *Acrosternum hilare*, green stink bug; *Euschistus servus*, brown stink bug; *Delia platura*, seedcorn maggot; *Mayetiola destructor*, Hessian fly; *Petrobia latens*, brown wheat mite; Oil Seed Rape: *Brevicoryne brassicae*, cabbage aphid; *Phyllotreta cruciferae*, Flea beetle; *Mamestra configurata*, Bertha armyworm; *Plutella xylostella*, Diamond-back moth; *Delia* spp., Root maggots.

10 [0131] In certain embodiments, the plant pathogen is selected from fungi, especially soil borne fungi such as *Fusarium oxysporum*, *Verticillium dahliae*, *Cladosporium* and *Ralstonia Solanaceum*.

3. Polynucleotides of the invention

[0132] The present invention is predicated, in part, on the isolation of two novel R 15 genes from banana, one from *Musa acuminata* Calcutta 4 and the other from *Musa acuminata* spp *malaccensis*. The Calcutta 4 gene designated RGA5 is 4380 nts long and comprises a single open reading frame of 4321 nts that encodes a 1441-aa putative polypeptide product. The nucleotide sequence of this gene and its deduced polypeptide sequence are presented in SEQ ID NO: 1 and 2, respectively. The *Musa acuminata* spp 20 *malaccensis* gene designated RGA2 comprises a single open reading frame of 3699 nts, which encodes a putative polypeptide product of 1232 aa. The nucleotide sequence of the RGA2 gene and its deduced polypeptide sequence are presented in SEQ ID NO: 3 and 4, respectively.

[0133] In accordance with the present invention, the novel R genes will be useful 25 for facilitating the construction of crop plants that are resistant to pathogenic disease, especially disease caused by fungal pathogens, viruses, nematodes, insects and the like. The R genes of the present invention can also be used as markers in genetic mapping as well as in assessing disease resistance in a plant of interest. Thus, the sequences can be used in breeding programs. See, for example, Gentzbittel *et al.* (1998, *Theor. Appl.* 30 *Genet.* 96:519-523). Additional uses for the sequences of the invention include using the sequences as bait to isolate other signalling components on defence/resistance pathways and to isolate the corresponding promoter sequences. The sequences may also

be used to modulate plant development processes, such as pollen development, regulation of organ shape, differentiation of aleurone and shoot epidermis, embryogenic competence, and cell/cell interactions. See, generally, Sambrook *et al.* (1989) Molecular Cloning: A Laboratory Manual (2nd ed., Cold Spring Harbor Laboratory Press, Plainview, N.Y.). The sequences of the present invention can also be used to generate variants (e.g., by 'domain swapping') for the generation of new resistance specificities.

[0134] The invention encompasses isolated or substantially purified nucleic acid or protein compositions. An "isolated" or "purified" nucleic acid molecule or protein, or biologically active portion thereof, is substantially or essentially free from components that normally accompany or interact with the nucleic acid molecule or protein as found in its naturally occurring environment. Thus, an isolated or purified polynucleotide or polypeptide is substantially free of other cellular material, or culture medium when produced by recombinant techniques, or substantially free of chemical precursors or other chemicals when chemically synthesised. Suitably, an "isolated" polynucleotide is free of sequences (especially protein encoding sequences) that naturally flank the polynucleotide (i.e., sequences located at the 5' and 3' ends of the polynucleotide) in the genomic DNA of the organism from which the polynucleotide was derived. For example, in various embodiments, the isolated polynucleotide can contain less than about 5 kb, 4 kb, 3 kb, 2 kb, 1 kb, 0.5 kb, or 0.1 kb of nucleotide sequences that naturally flank the polynucleotide in genomic DNA of the cell from which the polynucleotide was derived. A polypeptide that is substantially free of cellular material includes preparations of protein having less than about 30%, 20%, 10%, 5%, (by dry weight) of contaminating protein. When the protein of the invention or biologically active portion thereof is recombinantly produced, culture medium suitably represents less than about 30%, 20%, 10%, or 5% (by dry weight) of chemical precursors or non-protein-of-interest chemicals.

[0135] The present invention also encompasses portions of the disclosed nucleotide sequences. Portions of a nucleotide sequence may encode polypeptide portions or segments that retain the biological activity of the native polypeptide and hence modulate or regulate disease resistance. Alternatively, portions of a nucleotide sequence that are useful as hybridisation probes generally do not encode amino acid sequences retaining such biological activity. Thus, portions of a nucleotide sequence may range from at least about 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 40, 50,

60, 80, 90, 100 nucleotides, or almost up to the full-length nucleotide sequence encoding the polypeptides of the invention.

[0136] A portion of an R nucleotide sequence that encodes a biologically active portion of an R polypeptide of the invention will encode at least about 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 30, 40, 50, 60, 70, 80, 90, 100, 120, 150, 300, 400, 500, 600, 700, 800, 900 or 1000 contiguous amino acid residues, or almost up to the total number of amino acids present in a full-length R polypeptide of the invention (for example, 1440 or 1330 amino acid residues for SEQ ID NO: 2 or 4, respectively). Portions of an R nucleotide sequence that are useful as hybridisation probes or PCR primers generally need not encode a biologically active portion of an R polypeptide.

[0137] Thus, a portion of an R nucleotide sequence may encode a biologically active portion of an R polypeptide, or it may be a fragment that can be used as a hybridisation probe or PCR primer using standard methods known in the art. A biologically active portion of an R polypeptide can be prepared by isolating a portion of one of the R nucleotide sequences of the invention, expressing the encoded portion of the R polypeptide (e.g., by recombinant expression *in vitro*), and assessing the activity of the encoded portion of the R polypeptide. Nucleic acid molecules that are portions of an R nucleotide sequence comprise at least about 15, 16, 17, 18, 19, 20, 25, 30, 50, 75, 100, 150, 200, 250, 300, 350, 400, 450, 500, 550, 600, or 650 nucleotides, or almost up to the number of nucleotides present in a full-length R nucleotide sequence disclosed herein (for example, 4375 or 3690 nucleotides for SEQ ID NO: 1 or 14, respectively).

[0138] The invention also contemplates variants of the disclosed nucleotide sequences. Nucleic acid variants can be naturally occurring, such as allelic variants (same locus), homologues (different locus), and orthologues (different organism) or can be non naturally occurring. Naturally occurring variants such as these can be identified with the use of well-known molecular biology techniques, as, for example, with polymerase chain reaction (PCR) and hybridisation techniques as known in the art. Non-naturally occurring variants can be made by mutagenesis techniques, including those applied to polynucleotides, cells, or organisms. The variants can contain nucleotide substitutions, deletions, inversions and insertions. Variation can occur in either or both the coding and non-coding regions. The variations can produce both conservative and non-conservative amino acid substitutions (as compared in the encoded product). For nucleotide sequences,

conservative variants include those sequences that, because of the degeneracy of the genetic code, encode the amino acid sequence of one of the R polypeptides of the invention. Variant nucleotide sequences also include synthetically derived nucleotide sequences, such as those generated, for example, by using site-directed mutagenesis but
5 which still encode an R polypeptide of the invention. Generally, variants of a particular nucleotide sequence of the invention will have at least about 30%, 40% 50%, 55%, 60%, 65%, 70%, generally at least about 75%, 80%, 85%, desirably about 90% to 95% or more, and more suitably about 98% or more sequence identity to that particular nucleotide sequence as determined by sequence alignment programs described
10 elsewhere herein using default parameters.

[0139] The nucleotide sequences of the invention can be used to isolate corresponding sequences from other organisms, particularly other plants. Methods are readily available in the art for the hybridisation of nucleic acid sequences. Coding sequences from other plants may be isolated according to well known techniques based
15 on their sequence identity with the coding sequences set forth herein. In these techniques all or part of the known coding sequence is used as a probe which selectively hybridises to other R coding sequences present in a population of cloned genomic DNA fragments or cDNA fragments (i.e., genomic or cDNA libraries) from a chosen organism. Accordingly, the present invention also contemplates polynucleotides that
20 hybridise to the R gene nucleotide sequences, or to their complements, under stringency conditions described below. As used herein, the term "hybridises under low stringency, medium stringency, high stringency, or very high stringency conditions" describes conditions for hybridisation and washing. Guidance for performing hybridisation reactions can be found in Ausubel *et al.*, (1998, *supra*), Sections 6.3.1-6.3.6. Aqueous
25 and non-aqueous methods are described in that reference and either can be used. Reference herein to low stringency conditions include and encompass from at least about 1% v/v to at least about 15% v/v formamide and from at least about 1 M to at least about 2 M salt for hybridisation at 42° C, and at least about 1 M to at least about 2 M salt for washing at 42° C. Low stringency conditions also may include 1% Bovine
30 Serum Albumin (BSA), 1 mM EDTA, 0.5 M NaHPO₄ (pH 7.2), 7% SDS for hybridisation at 65° C, and (i) 2xSSC, 0.1% SDS; or (ii) 0.5% BSA, 1 mM EDTA, 40 mM NaHPO₄ (pH 7.2), 5% SDS for washing at room temperature. One embodiment of low stringency conditions includes hybridisation in 6X sodium chloride/sodium citrate (SSC) at about 45° C, followed by two washes in 0.2X SSC, 0.1% SDS at least at 50° C

(the temperature of the washes can be increased to 55° C for low stringency conditions). Medium stringency conditions include and encompass from at least about 16% v/v to at least about 30% v/v formamide and from at least about 0.5 M to at least about 0.9 M salt for hybridisation at 42° C, and at least about 0.1 M to at least about 0.2 M salt for washing at 55° C. Medium stringency conditions also may include 1% Bovine Serum Albumin (BSA), 1 mM EDTA, 0.5 M NaHPO₄ (pH 7.2), 7% SDS for hybridisation at 65° C, and (i) 2 x SSC, 0.1% SDS; or (ii) 0.5% BSA, 1 mM EDTA, 40 mM NaHPO₄ (pH 7.2), 5% SDS for washing at 60-65° C. One embodiment of medium stringency conditions includes hybridising in 6X SSC at about 45° C, followed by one or more washes in 0.2X SSC, 0.1% SDS at 60° C. High stringency conditions include and encompass from at least about 31% v/v to at least about 50% v/v formamide and from about 0.01 M to about 0.15 M salt for hybridisation at 42° C, and about 0.01 M to about 0.02 M salt for washing at 55° C. High stringency conditions also may include 1% BSA, 1 mM EDTA, 0.5 M NaHPO₄ (pH 7.2), 7% SDS for hybridisation at 65° C, and (i) 0.2 x SSC, 0.1% SDS; or (ii) 0.5% BSA, 1mM EDTA, 40 mM NaHPO₄ (pH 7.2), 1% SDS for washing at a temperature in excess of 65° C. One embodiment of high stringency conditions includes hybridising in 6X SSC at about 45° C, followed by one or more washes in 0.2X SSC, 0.1% SDS at 65° C.

[0140] In certain embodiments, an isolated nucleic acid molecule of the invention hybridises under very high stringency conditions. One embodiment of very high stringency conditions includes hybridising 0.5M sodium phosphate, 7% SDS at 65°C, followed by one or more washes at 0.2X SSC, 1% SDS at 65° C.

[0141] Other stringency conditions are well known in the art and a skilled addressee will recognise that various factors can be manipulated to optimise the specificity of the hybridisation. Optimisation of the stringency of the final washes can serve to ensure a high degree of hybridisation. For detailed examples, see Ausubel *et al.*, *supra* at pages 2.10.1 to 2.10.16 and Sambrook *et al.* (1989, *supra*) at sections 1.101 to 1.104.

[0142] While stringent washes are typically carried out at temperatures from about 42° C to 68° C, one skilled in the art will appreciate that other temperatures may be suitable for stringent conditions. Maximum hybridisation rate typically occurs at about 20° C to 25° C below the T_m for formation of a DNA-DNA hybrid. It is well known in the art that the T_m is the melting temperature, or temperature at which two

complementary polynucleotide sequences dissociate. Methods for estimating T_m are well known in the art (see Ausubel *et al.*, *supra* at page 2.10.8). In general, the T_m of a perfectly matched duplex of DNA may be predicted as an approximation by the formula:

5
$$T_m = 81.5 + 16.6 (\log_{10} M) + 0.41 (\%G+C) - 0.63 (\% \text{ formamide}) - (600/\text{length})$$

wherein: M is the concentration of Na^+ , preferably in the range of 0.01 molar to 0.4 molar; %G+C is the sum of guanosine and cytosine bases as a percentage of the total number of bases, within the range between 30% and 75% G+C; % formamide is the percent formamide concentration by volume; length is the number of base pairs in the
10 DNA duplex. The T_m of a duplex DNA decreases by approximately 1° C with every increase of 1% in the number of randomly mismatched base pairs. Washing is generally carried out at $T_m - 15^\circ \text{C}$ for high stringency, or $T_m - 30^\circ \text{C}$ for moderate stringency.

[0143] In one example of a hybridisation procedure, a membrane (e.g., a nitrocellulose membrane or a nylon membrane) containing immobilised DNA is
15 hybridised overnight at 42° C in a hybridisation buffer (50% deionised formamide, 5xSSC, 5x Denhardt's solution (0.1% ficoll, 0.1% polyvinylpyrrolidone and 0.1% bovine serum albumin), 0.1% SDS and 200 mg/mL denatured salmon sperm DNA) containing labelled probe. The membrane is then subjected to two sequential medium stringency washes (i.e., 2xSSC, 0.1% SDS for 15 min at 45° C, followed by 2xSSC,
20 0.1% SDS for 15 min at 50° C), followed by two sequential higher stringency washes (i.e., 0.2xSSC, 0.1% SDS for 12 min at 55° C followed by 0.2xSSC and 0.1%SDS solution for 12 min at 65-68° C).

[0144] Variant nucleotide sequences also encompass sequences derived from a mutagenic or recombinogenic procedures such as 'DNA shuffling' which can be used
25 for swapping domains in a polypeptide of interest with domains of other polypeptides. With DNA shuffling, one or more different R coding sequences can be manipulated to create a new R sequence possessing desired properties. In this procedure, libraries of recombinant polynucleotides are generated from a population of related polynucleotides comprising sequence regions that have substantial sequence identity and can be
30 homologously recombined *in vitro* or *in vivo*. For example, using this approach, sequence motifs encoding a domain of interest (e.g., the coiled coil domain, the NBS domain and/or the LRR domain of the polypeptides of the invention) may be shuffled between the R gene of the invention and other known R genes to obtain a new gene

coding for a protein with an improved property of interest, such broadening spectrum of disease resistance. Illustrative resistance R genes that could be employed for this purpose are listed in Table B below.

TABLE B

5

PLANT DISEASE RESISTANCE GENES CLONED FROM 1994 TO 2003.

PLANT-PATHOGEN INTERACTION		PLANT (R) PROTEIN STRUCTURE	R PROTEIN NAME	REFERENCE
Flax	<i>Melampsora lini</i>	TIR-NBS-LRR	L	Lawrence <i>et al.</i> , 1995
TOBACCO	Tobacco mosaic virus	TIR-NBS-LRR	N	Whitman <i>et al.</i> , 1996
Flax	<i>Melampsora lini</i>	TIR-NBS-LRR	M	Anderson <i>et al.</i> , 1997
ARABIDOPSIS	<i>Peronospora parasitica</i>	TIR-NBS-LRR	RPP 5	Parker <i>et al.</i> , 1997
Arabidopsis	<i>Pseudomonas</i>	TIR-NBS-LRR	RPS4	Gassmann <i>et al.</i> , 1999
Flax	<i>Melampsora lini</i>	TIR-NBS-LRR	P	Dodds <i>et al.</i> , 2000
3.1.1 Arabidopsis	<i>Pseudomonas syringae</i>	NBS-LRR	RPS2	Mindrinis <i>et al.</i> , 1994
	<i>Pseudomonas syringae</i>	NBS-LRR	Prf	Salmeron <i>et al.</i> , 1996
Tomato	<i>Pseudomonas syringae</i>	NBS-LRR	RPM1	Grant <i>et al.</i> , 1996
Wheat	<i>Heterodera avenae</i>	NBS-LRR	Cre3	Lagudah <i>et al.</i> , 1997
Tomato	3.2 <i>Fusarium</i>	NBS-LRR	12	Simons <i>et al.</i> , 1998
Tomato	3.3 <i>Meloidogyne</i> sp.	NBS-LRR	Mi	Milligan <i>et al.</i> , 1998
Tomato	<i>Macrosiphum euphorbiae</i>	NBS-LRR	Mi	Milligan <i>et al.</i> , 1998
Arabidopsis	<i>Peronospora parasitica</i>	NBS-LRR	RPP1	Botella <i>et al.</i> , 1998
Lettuce	<i>Bremia lactucae</i>	NBS-LRR	Dm3	Meyers <i>et al.</i> , 1998
Rice	<i>Xanthomonas</i>	NBS-LRR	Xa1	Yoshimura <i>et al.</i> , 1998
Arabidopsis	<i>Pseudomonas</i>	NBS-LRR	RPS5	Warren <i>et al.</i> , 1998
Maize	<i>Puccinia sorghi</i>	NBS-LRR	Rp1-D	Collins <i>et al.</i> , 1999
Pepper	<i>Xanthomonas campestris</i>	NBS-LRR	Bs2	Thai <i>et al.</i> , 1999
Potato	PVX	NBS-LRR	Rx2	Bendahmane <i>et al.</i> , 1999
Rice	<i>Magnaporthe</i>	NBS-LRR	Pi-ta	Bryan <i>et al.</i> , 2000
Barley	<i>Blumeria graminis</i>	NBS-LRR	Mla	Zhou <i>et al.</i> , 2000
Arabidopsis	<i>Peronospora parasitica</i>	NBS-LRR	RPP 13	Bitner-Eddy <i>et al.</i> , 2000
Tomato	Tospovirus	NBS-LRR	Sw-5	Brommonschenkel <i>et al.</i> , 2000

PLANT-PATHOGEN INTERACTION		PLANT (R) PROTEIN STRUCTURE	R PROTEIN NAME	REFERENCE
Potato	<i>Globodera pallida</i>	NBS-LRR	Gpa 2	Van der Vossen <i>et al.</i> , 2000
Potato	<i>Phytophthora infestans</i>	NBS-LRR	R1	Ballvora <i>et al.</i> , 2002
Tomato	<i>Globodera rostochiensis</i>	NBS-LRR	Hero	Ernst <i>et al.</i> , 2002
Potato	<i>Phytophthora infestans</i>	NBS-LRR	RB	Song <i>et al.</i> , 2003

[0145] Strategies for DNA shuffling are known in the art. See, for example: Stemmer (1994, *Proc. Natl. Acad. Sci. USA* 91:10747-10751; 1994, *Nature* 370:389-391); Crameri *et al.* (1997, *Nature Biotech.* 15:436-438); Moore *et al.* (1997, *J. Mol. Biol.* 272:336-347); Zhang *et al.* (1997 *Proc. Natl. Acad. Sci. USA* 94:4504-4509);
5 Crameri *et al.* (1998, *Nature* 391:288-291); and U.S. Pat. Nos. 5,605,793 and 5,837,458.

4. Polypeptides of the invention

[0146] The present invention provides polypeptides and biologically active portions thereof that confer resistance to disease, especially resistance to pathogenic disease
10 including disease caused by fungal pathogen, viruses, nematodes, insects and the like. Biologically active portions of the R polypeptides of the invention include portions with immuno-interactive activity of at least about 6, 8, 10, 12, 14, 16, 18, 20, 25, 30, 40, 50, 60 amino acid residues in length. For example, immuno-interactive fragments contemplated by the present invention are at least 6 and desirably at least 8 amino acid
15 residues in length, which can elicit an immune response in an animal for the production of antigen-binding molecules that are immuno-interactive with the R polypeptides of the invention. Such antigen-binding molecules can be used to screen organisms, especially plants, for structurally and/or functionally related R polypeptides. Typically, portions of the disclosed R polypeptides may participate in an interaction, e.g., an intramolecular or
20 an inter-molecular interaction. An inter-molecular interaction can be a specific binding interaction or an enzymatic interaction (e.g., the interaction can be transient and a covalent bond is formed or broken). An inter-molecular interaction can be between an R polypeptide and a pathogen elicitor protein. Biologically active portions of an R polypeptide include peptides comprising amino acid sequences sufficiently similar to or
25 derived from the amino acid sequences of the disclosed R polypeptides, e.g., the amino acid sequences shown in SEQ ID NO: 2 or 4, which include less amino acids than the full-length R polypeptide, and exhibit at least one activity of an R polypeptide.

Typically, biologically active portions comprise a domain or motif with at least one activity of the R polypeptide, e.g., the ability to bind to a pathogen elicitor protein or to confer disease resistance. A biologically active portion of an R polypeptide can be a polypeptide which is, for example, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 30, 40, 50, 60, 70, 80, 90, 100, 120, 150, 300, 400, 500, 600, 700, 800, 900 or 1000 or more amino acids in length. Suitably, the portion is a "biologically-active portion" having no less than about 1%, 10%, 25% 50% of the pathogen elicitor protein-binding activity or the resistance-conferring activity of the full-length polypeptide.

10 [0147] The present invention also contemplates variant R polypeptides. "Variant" polypeptides include proteins derived from the native protein by deletion (so-called truncation) or addition of one or more amino acids to the N-terminal and/or C-terminal end of the native protein; deletion or addition of one or more amino acids at one or more sites in the native protein; or substitution of one or more amino acids at one or more sites in the native protein. Variant proteins encompassed by the present invention are biologically active, that is they continue to possess the desired biological activity of the native protein, that is, modulating disease resistance or interacting with a pathogen elicitor protein. Such variants may result from, for example, genetic polymorphism or from human manipulation. Biologically active variants of a native R protein of the invention will have at least 40%, 50%, 60%, 70%, generally at least 75%, 80%, 85%, preferably about 90% to 95% or more, and more preferably about 98% or more sequence similarity with the amino acid sequence for the native protein as determined by sequence alignment programs described elsewhere herein using default parameters. A biologically active variant of a protein of the invention may differ from that protein by as few as 1-15 amino acid residues, as few as 1-10, such as 6-10, as few as 5, as few as 4, 3, 2, or even 1 amino acid residue.

[0148] The proteins of the invention may be altered in various ways including amino acid substitutions, deletions, truncations, and insertions. Methods for such manipulations are generally known in the art. For example, amino acid sequence variants of the R proteins can be prepared by mutations in the DNA. Methods for mutagenesis and nucleotide sequence alterations are well known in the art. See, for example, Kunkel (1985, *Proc. Natl. Acad. Sci. USA* 82:488-492), Kunkel *et al.* (1987, *Methods in Enzymol.* 154:367-382), U.S. Pat. No. 4,873,192, Watson, J. D. *et al.*

("Molecular Biology of the Gene", Fourth Edition, Benjamin/Cummings, Menlo Park, Calif., 1987) and the references cited therein. Guidance as to appropriate amino acid substitutions that do not affect biological activity of the protein of interest may be found in the model of Dayhoff et al. (1978) Atlas of Protein Sequence and Structure (Natl. Biomed. Res. Found., Washington, D.C.). Methods for screening gene products of combinatorial libraries made by point mutations or truncation, and for screening cDNA libraries for gene products having a selected property are known in the art. Such methods are adaptable for rapid screening of the gene libraries generated by combinatorial mutagenesis of R polypeptides. Recursive ensemble mutagenesis (REM), a new technique which enhances the frequency of functional mutants in the libraries, can be used in combination with the screening assays to identify R polypeptide variants (Arkin and Yourvan (1992) *Proc. Natl. Acad. Sci. USA* 89:7811-7815; Delgrave et al. (1993) *Protein Engineering* 6:327-331). Conservative substitutions, such as exchanging one amino acid with another having similar properties, may be desirable as discussed in more detail below.

[0149] Variant polypeptides may contain conservative amino acid substitutions at various locations along their sequence, as compared to the R polypeptide amino acid sequences of the invention. A "conservative amino acid substitution" is one in which the amino acid residue is replaced with an amino acid residue having a similar side chain. Families of amino acid residues having similar side chains have been defined in the art, which can be generally sub-classified as follows:

[0150] Acidic: The residue has a negative charge due to loss of H ion at physiological pH and the residue is attracted by aqueous solution so as to seek the surface positions in the conformation of a peptide in which it is contained when the peptide is in aqueous medium at physiological pH. Amino acids having an acidic side chain include glutamic acid and aspartic acid.

[0151] Basic: The residue has a positive charge due to association with H ion at physiological pH or within one or two pH units thereof (e.g., histidine) and the residue is attracted by aqueous solution so as to seek the surface positions in the conformation of a peptide in which it is contained when the peptide is in aqueous medium at physiological pH. Amino acids having a basic side chain include arginine, lysine and histidine.

[0152] Charged: The residues are charged at physiological pH and, therefore, include amino acids having acidic or basic side chains (i.e., glutamic acid, aspartic acid, arginine, lysine and histidine).

5 [0153] Hydrophobic: The residues are not charged at physiological pH and the residue is repelled by aqueous solution so as to seek the inner positions in the conformation of a peptide in which it is contained when the peptide is in aqueous medium. Amino acids having a hydrophobic side chain include tyrosine, valine, isoleucine, leucine, methionine, phenylalanine and tryptophan.

10 [0154] Neutral/polar: The residues are not charged at physiological pH, but the residue is not sufficiently repelled by aqueous solutions so that it would seek inner positions in the conformation of a peptide in which it is contained when the peptide is in aqueous medium. Amino acids having a neutral/polar side chain include asparagine, glutamine, cysteine, histidine, serine and threonine.

15 [0155] This description also characterises certain amino acids as "small" since their side chains are not sufficiently large, even if polar groups are lacking, to confer hydrophobicity. With the exception of proline, "small" amino acids are those with four carbons or less when at least one polar group is on the side chain and three carbons or less when not. Amino acids having a small side chain include glycine, serine, alanine and threonine. The gene-encoded secondary amino acid proline is a special case due to
20 its known effects on the secondary conformation of peptide chains. The structure of proline differs from all the other naturally-occurring amino acids in that its side chain is bonded to the nitrogen of the α -amino group, as well as the α -carbon. Several amino acid similarity matrices (e.g., PAM120 matrix and PAM250 matrix as disclosed for example by Dayhoff *et al.* (1978) A model of evolutionary change in proteins. Matrices
25 for determining distance relationships In M. O. Dayhoff, (ed.), Atlas of protein sequence and structure, Vol. 5, pp. 345-358, National Biomedical Research Foundation, Washington DC; and by Gonnet *et al.*, 1992, *Science* 256(5062): 144301445), however, include proline in the same group as glycine, serine, alanine and threonine. Accordingly, for the purposes of the present invention, proline is classified as a "small"
30 amino acid.

[0156] The degree of attraction or repulsion required for classification as polar or nonpolar is arbitrary and, therefore, amino acids specifically contemplated by the

invention have been classified as one or the other. Most amino acids not specifically named can be classified on the basis of known behaviour.

[0157] Amino acid residues can be further sub-classified as cyclic or noncyclic, and aromatic or nonaromatic, self-explanatory classifications with respect to the side-chain substituent groups of the residues, and as small or large. The residue is considered small if it contains a total of four carbon atoms or less, inclusive of the carboxyl carbon, provided an additional polar substituent is present; three or less if not. Small residues are, of course, always nonaromatic. Dependent on their structural properties, amino acid residues may fall in two or more classes. For the naturally-occurring protein amino acids, sub-classification according to the this scheme is presented in the Table C.

TABLE C

AMINO ACID SUB-CLASSIFICATION

Sub-classes	Amino acids
Acidic	Aspartic acid, Glutamic acid
Basic	Noncyclic: Arginine, Lysine; Cyclic: Histidine
Charged	Aspartic acid, Glutamic acid, Arginine, Lysine, Histidine
Small	Glycine, Serine, Alanine, Threonine, Proline
Polar/neutral	Asparagine, Histidine, Glutamine, Cysteine, Serine, Threonine
Polar/large	Asparagine, Glutamine
Hydrophobic	Tyrosine, Valine, Isoleucine, Leucine, Methionine, Phenylalanine, Tryptophan
Aromatic	Tryptophan, Tyrosine, Phenylalanine
Residues that influence chain orientation	Glycine and Proline

[0158] Accordingly, the present invention also contemplates variants of the naturally occurring or parent R polypeptide sequences or their biologically-active fragments, wherein the variants are distinguished from the parent sequences by the addition, deletion, or substitution of one or more amino acids. In general, variants display at least about 30, 40, 50, 55, 60, 65, 70, 75, 80, 85, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99 % similarity to a parent R polypeptide sequence as for example set forth in SEQ

ID NO: 2 or 4. Desirably, variants will have at least 30, 40, 50, 55, 60, 65, 70, 75, 80, 85, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99 % sequence identity to a parent R polypeptide sequence as set forth in SEQ ID NO:2 or 4. Moreover, sequences differing from the native or parent sequences by the addition, deletion, or substitution of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 30, 40, 50, 60, 70, 80, 90, 100, 150, 200, 300, 500 or more amino acids but which retain the disease-resistance-conferring or pathogen elicitor-binding properties are contemplated. Polypeptides of the invention include polypeptides that are encoded by polynucleotides that hybridise under stringency conditions as defined herein, especially high stringency conditions, to the polynucleotide sequences of the invention, or the non-coding strand thereof, as described above.

[0159] In one embodiment, variant polypeptides differ from the disclosed sequences by at least one but by less than 50, 40, 30, 20, 15, 10, 8, 6, 5, 4, 3, 2 or 1 amino acid residue(s). In another, variant polypeptides differ from the corresponding sequence in SEQ ID NO: 2 or 4 by at least 1% but less than 20%, 15%, 10% or 5% of the residues. (If this comparison requires alignment the sequences should be aligned for maximum similarity. "Looped" out sequences from deletions or insertions, or mismatches, are considered differences.) The differences are, suitably, differences or changes at a non-essential residue or a conservative substitution.

[0160] A "non-essential" amino acid residue is a residue that can be altered from the wild-type sequence of an R polypeptide without abolishing or substantially altering one or more of its activities (e.g., disease-resistance or pathogen elicitor-binding properties). Suitably, the alteration does not substantially alter one of these activities, e.g., the activity is at least 20%, 40%, 60%, 70% or 80% of wild-type. An "essential" amino acid residue is a residue that, when altered from the wild-type sequence of an R polypeptide of the invention, results in abolition of disease-resistance or pathogen elicitor-binding properties such that less than 20% of the wild-type activity is present. For example, conserved amino acid residues between the R polypeptides shown in Figure 2 may be unamenable to alteration.

[0161] Desirable variant R polypeptides are those having conserved amino acid substitutions. Examples of conservative substitutions include the following: aspartic-glutamic as acidic amino acids; lysine/arginine/histidine as basic amino acids; serine/glycine/alanine/threonine as small amino acids; leucine/isoleucine,

methionine/valine, alanine/valine as hydrophobic amino acids. Conservative amino acid substitution also includes groupings based on side chains. For example, a group of amino acids having aliphatic side chains is glycine, alanine, valine, leucine, and isoleucine; a group of amino acids having aliphatic-hydroxyl side chains is serine and threonine; a group of amino acids having amide-containing side chains is asparagine and glutamine; a group of amino acids having aromatic side chains is phenylalanine, tyrosine, and tryptophan; a group of amino acids having basic side chains is lysine, arginine, and histidine; and a group of amino acids having sulphur-containing side chains is cysteine and methionine. For example, it is reasonable to expect that replacement of a leucine with an isoleucine or valine, an aspartate with a glutamate, a threonine with a serine, or a similar replacement of an amino acid with a structurally related amino acid will not have a major effect on the properties of the resulting variant polypeptide. Whether an amino acid change results in a functional R polypeptide can readily be determined by assaying its disease resistance-conferring activity or its pathogen-elicitor-binding activity. Conservative substitutions are shown in Table D below under the heading of exemplary substitutions. More preferred substitutions are shown under the heading of preferred substitutions. Amino acid substitutions falling within the scope of the invention, are, in general, accomplished by selecting substitutions that do not differ significantly in their effect on maintaining (a) the structure of the peptide backbone in the area of the substitution, (b) the charge or hydrophobicity of the molecule at the target site, or (c) the bulk of the side chain. After the substitutions are introduced, the variants are screened for biological activity.

TABLE D**EXEMPLARY AND PREFERRED AMINO ACID SUBSTITUTIONS**

<i>Original Residue</i>	Exemplary Substitutions	Preferred Substitutions
Ala	Val, Leu, Ile	Val
Arg	Lys, Gln, Asn	Lys
Asn	Gln, His, Lys, Arg	Gln
Asp	Glu	Glu
Cys	Ser	Ser
Gln	Asn, His, Lys,	Asn

<i>Original Residue</i>	<i>Exemplary Substitutions</i>	<i>Preferred Substitutions</i>
Glu	Asp, Lys	Asp
Gly	Pro	Pro
His	Asn, Gln, Lys, Arg	Arg
Ile	Leu, Val, Met, Ala, Phe, Norleu	Leu
Leu	Norleu, Ile, Val, Met, Ala, Phe	Ile
Lys	Arg, Gln, Asn	Arg
Met	Leu, Ile, Phe	Leu
Phe	Leu, Val, Ile, Ala	Leu
Pro	Gly	Gly
Ser	Thr	Thr
Thr	Ser	Ser
Trp	Tyr	Tyr
Tyr	Trp, Phe, Thr, Ser	Phe
Val	Ile, Leu, Met, Phe, Ala, Norleu	Leu

[0162] Alternatively, similar amino acids for making conservative substitutions can be grouped into three categories based on the identity of the side chains. The first group includes glutamic acid, aspartic acid, arginine, lysine, histidine, which all have charged side chains; the second group includes glycine, serine, threonine, cysteine, tyrosine, glutamine, asparagine; and the third group includes leucine, isoleucine, valine, alanine, proline, phenylalanine, tryptophan, methionine, as described in Zubay, G., *Biochemistry*, third edition, Wm.C. Brown Publishers (1993).

[0163] Thus, a predicted non-essential amino acid residue in an R polypeptide is typically replaced with another amino acid residue from the same side chain family. Alternatively, mutations can be introduced randomly along all or part of an R gene coding sequence, such as by saturation mutagenesis, and the resultant mutants can be screened for disease resistance-conferring activity or pathogen-elicitor-binding activity to identify mutants that retain activity. Following mutagenesis of the coding sequences,

the encoded peptide can be expressed recombinantly and the activity of the peptide can be determined.

[0164] In other embodiments, the SCE includes an amino acid sequence having at least about 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%
5 95%, 96%, 97%, 98% or more similarity to a corresponding sequence of SEQ ID NO: 2 or 4, and has disease resistance-conferring activity or pathogen-elicitor-binding activity.

[0165] The R polypeptides of the present invention contain a significant number of structural characteristics in common with each other as for example depicted in Figure 2. The term "family" when referring to the protein and nucleic acid molecules of the
10 invention means two or more proteins or nucleic acid molecules having a common structural domain or motif and having sufficient amino acid or nucleotide sequence homology as defined herein. Such family members can be naturally or non-naturally-occurring and can be from either the same or different species. Members of a family can also have common functional characteristics.

15 **5. *Anti-R polypeptide antigen-binding molecules***

[0166] The invention also provides an antigen-binding molecule that is specifically immuno-interactive with an R polypeptide of the invention. In one embodiment, the antigen-binding molecule comprise whole polyclonal antibodies. Such antibodies may be prepared, for example, by injecting a polypeptide, portion or variant of the invention
20 into a production species, which may include mice or rabbits, to obtain polyclonal antisera. Methods of producing polyclonal antibodies are well known to those skilled in the art. Exemplary protocols which may be used are described for example in Coligan *et al.*, CURRENT PROTOCOLS IN IMMUNOLOGY, (John Wiley & Sons, Inc, 1991), and Ausubel *et al.*, (1994-1998, *supra*), in particular Section III of Chapter 11.

25 [0167] In lieu of polyclonal antisera obtained in a production species, monoclonal antibodies may be produced using the standard method as described, for example, by Köhler and Milstein (1975, *Nature* 256, 495-497), or by more recent modifications thereof as described, for example, in Coligan *et al.*, (1991, *supra*) by immortalising spleen or other antibody producing cells derived from a production species which has
30 been inoculated with one or more of the polypeptides, fragments, variants or derivatives of the invention.

- [0168] The invention also contemplates as antigen-binding molecules Fv, Fab, Fab' and F(ab')₂ immunoglobulin fragments. Alternatively, the antigen-binding molecule may comprise a synthetic stabilised Fv fragment. Exemplary fragments of this type include single chain Fv fragments (sFv, frequently termed scFv) in which a peptide linker is used to bridge the N terminus or C terminus of a V_H domain with the C terminus or N-terminus, respectively, of a V_L domain. ScFvs lack all constant parts of whole antibodies and are not able to activate complement. ScFvs may be prepared, for example, in accordance with methods outlined in Kreber *et al.* (Kreber *et al.* 1997, *J. Immunol. Methods*; 201(1): 35-55). Alternatively, they may be prepared by methods described in U.S. Patent No 5,091,513, European Patent No 239,400 or the articles by Winter and Milstein (1991, *Nature* 349:293) and Plückthun *et al.* (1996, In *Antibody engineering: A practical approach*. 203-252). In another embodiment, the synthetic stabilised Fv fragment comprises a disulphide stabilised Fv (dsFv) in which cysteine residues are introduced into the V_H and V_L domains such that in the fully folded Fv molecule the two residues will form a disulphide bond therebetween. Suitable methods of producing dsFv are described for example in (Glockscuther *et al.* *Biochem.* 29: 1363-1367; Reiter *et al.* 1994, *J. Biol. Chem.* 269: 18327-18331; Reiter *et al.* 1994, *Biochem.* 33: 5451-5459; Reiter *et al.* 1994, *Cancer Res.* 54: 2714-2718; Webber *et al.* 1995, *Mol. Immunol.* 32: 249-258).
- [0169] Phage display and combinatorial methods for generating R polypeptide antigen-binding molecules are known in the art (as described in, e.g., Ladner *et al.* U.S. Patent No. 5,223,409; Kang *et al.* International Publication No. WO 92/18619; Dower *et al.* International Publication No. WO 91/17271; Winter *et al.* International Publication WO 92/20791; Markland *et al.* International Publication No. WO 92/15679; Breitling *et al.* International Publication WO 93/01288; McCafferty *et al.* International Publication No. WO 92/01047; Garrard *et al.* International Publication No. WO 92/09690; Ladner *et al.* International Publication No. WO 90/02809; Fuchs *et al.* (1991) *Bio/Technology* 9:1370-1372; Hay *et al.* (1992) *Hum Antibod Hybridomas* 3:81-85; Huse *et al.* (1989) *Science* 246:1275-1281; Griffiths *et al.* (1993) *EMBO J* 12:725-734; Hawkins *et al.* (1992) *J Mol Biol* 226:889-896; Clackson *et al.* (1991) *Nature* 352:624-628; Gram *et al.* (1992) *PNAS* 89:3576-3580; Garrad *et al.* (1991) *Bio/Technology* 9:1373-1377; Hoogenboom *et al.* (1991) *Nuc Acid Res* 19:4133-4137; and Barbas *et al.* (1991) *PNAS* 88:7978-7982). The antigen-binding molecules can be used to screen expression libraries for variant R polypeptides. They can also be used to detect and/or isolate the R

polypeptides of the invention. Thus, the invention also contemplates the use of antigen-binding molecules to isolate R polypeptides using , for example, any suitable immunoaffinity based method including, but not limited to, immunochromatography and immunoprecipitation. A suitable method utilises solid phase adsorption in which
5 anti-R polypeptide antigen-binding molecules are attached to a suitable resin, the resin is contacted with a sample suspected of containing a natriuretic peptides, and the natriuretic peptides, if any, are subsequently eluted from the resin. Illustrative resins include: Sepharose® (Pharmacia), Poros® resins (Roche Molecular Biochemicals, Indianapolis), Actigel Superflow™ resins (Sterogene Bioseparations Inc., Carlsbad
10 Calif.), and Dynabeads™ (Dynal Inc., Lake Success, N.Y.).

[0170] The antigen-binding molecule can be coupled to a compound, e.g., a label such as a radioactive nucleus, or imaging agent, e.g. a radioactive, enzymatic, or other, e.g., imaging agent, e.g., a NMR contrast agent. Labels which produce detectable radioactive emissions or fluorescence are preferred. An anti-R polypeptide antigen-
15 binding molecule (e.g., monoclonal antibody) can be used to detect R polypeptides (e.g., in a cellular lysate or cell supernatant) in order to evaluate the abundance and pattern of expression of the protein. Anti-R polypeptides antigen-binding molecules can be used diagnostically to monitor R polypeptides levels in tissue as part of an agronomic testing procedure. Detection can be facilitated by coupling (i.e., physically
20 linking) the antibody to a detectable substance (i.e., antibody labelling). Examples of detectable substances include various enzymes, prosthetic groups, fluorescent materials, luminescent materials, bioluminescent materials, and radioactive materials. Examples of suitable enzymes include horseradish peroxidase, alkaline phosphatase, β -galactosidase, or acetylcholinesterase; examples of suitable prosthetic group complexes include
25 streptavidin/biotin and avidin/biotin; examples of suitable fluorescent materials include umbelliferone, fluorescein, fluorescein isothiocyanate, rhodamine, dichlorotriazinylamine fluorescein, dansyl chloride or phycoerythrin; an example of a luminescent material includes luminol; examples of bioluminescent materials include luciferase, luciferin, and aequorin, and examples of suitable radioactive material include
30 ^{125}I , ^{131}I , ^{35}S or ^3H . The label may be selected from a group including a chromogen, a catalyst, an enzyme, a fluorophore, a chemiluminescent molecule, a lanthanide ion such as Europium (Eu^{34}), a radioisotope and a direct visual label. In the case of a direct visual label, use may be made of a colloidal metallic or non-metallic particle, a dye particle, an

enzyme or a substrate, an organic polymer, a latex particle, a liposome, or other vesicle containing a signal producing substance and the like.

[0171] A large number of enzymes useful as labels is disclosed in United States Patent Specifications U.S. 4,366,241, U.S. 4,843,000, and U.S. 4,849,338. Enzyme
5 labels useful in the present invention include alkaline phosphatase, horseradish peroxidase, luciferase, β -galactosidase, glucose oxidase, lysozyme, malate dehydrogenase and the like. The enzyme label may be used alone or in combination with a second enzyme in solution.

6. Nucleic acid constructs

10 6.1 Prokaryotic expression

[0172] The present invention further relates to a nucleic acid construct designed for genetic transformation of prokaryotic cells, comprising a polynucleotide, portion or variant according to the invention operably linked to a regulatory sequence, which will typically comprise a transcriptional control element or promoter. Suitably, the chimeric
15 construct is operable in a Gram-negative prokaryotic cell. A variety of prokaryotic expression vectors, which may be used as a basis for constructing the chimeric nucleic acid construct, may be utilised to express a polynucleotide, portion or variant according to the invention. These include but are not limited to a chromosomal vector (e.g., a bacteriophage such as bacteriophage λ), an extrachromosomal vector (e.g., a plasmid or
20 a cosmid expression vector). The expression vector will also typically contain an origin of replication, which allows autonomous replication of the vector, and one or more genes that allow phenotypic selection of the transformed cells. Any of a number of suitable promoter sequences, including constitutive and inducible promoter sequences, may be used in the expression vector (see e.g., Bitter, *et al.*, 1987, *Methods in*
25 *Enzymology* 153: 516-544). For example, inducible promoters such as pL of bacteriophage γ , plac, ptrp, ptac ptrp-lac hybrid promoter and the like may be used. The nucleic acid construct may then be used to transform the desired prokaryotic host cell to produce a recombinant prokaryotic host cell, e.g., for producing a recombinant R polypeptide.

30 6.2 Eukaryotic expression

[0173] The invention also contemplates a nucleic acid construct designed for expressing a polynucleotide, portion or variant of the invention in a eukaryotic host cell.

A variety of eukaryotic host-expression vector systems may be utilised in this regard. These include, but are not limited to, yeast transformed with recombinant yeast expression vectors; insect cell systems infected with recombinant virus expression vectors (e.g., baculovirus); or animal cell systems infected with recombinant virus expression vectors (e.g., retroviruses, adenovirus, Vaccinia virus), or transformed animal cell systems engineered for stable expression. In certain advantageous embodiments, the chimeric nucleic acid construct is designed for genetic transformation of plants as described hereinafter.

6.3 Plant expression

10 [0174] In accordance with the present invention, it is proposed that the R gene polynucleotides, portions and variants of the invention will be useful for facilitating the construction of crop plants that are resistant to pathogenic disease, including diseases caused by fungal pathogens, viruses, nematodes, insects and the like. Accordingly, the present invention also relates to operably linking a polynucleotide, portion or variant of
15 as described hereinto a regulatory sequence (e.g., a promoter and a 3' non-translated region) that is function in plants to create a nucleic acid construct, designed for genetic transformation of plants.

6.3.1 Plant promoters

[0175] Numerous promoters that are active in plant cells have been described in the
20 literature, illustrative examples of which include the nopaline synthase (NOS) promoter, the octopine synthase (OCS) promoter (which is carried on tumour-inducing plasmids of *Agrobacterium tumefaciens*), the caulimovirus promoters such as the cauliflower mosaic virus (CaMV) 19S promoter and the CaMV 35S promoter, the figwort mosaic virus 35S-promoter, the light-inducible promoter from the small subunit of ribulose-1,5-
25 bis-phosphate carboxylase (ssRUBISCO), the Adh promoter, the sucrose synthase promoter, the R gene complex promoter, the GST-II-27 gene promoter and the chlorophyll a/b binding protein gene promoter, etc.

[0176] For the purpose of expression in source tissues of the plant, such as the leaf, seed, root or stem, it is desirable that the promoters driving expression of the target gene
30 have relatively high expression in these specific tissues. For this purpose, one may choose from a number of promoters for genes with tissue- or cell-specific or enhanced expression. Examples of such promoters reported in the literature include the

chloroplast glutamine synthetase GS2 promoter from pea, the chloroplast fructose-1,6-biphosphatase (FBPase) promoter from wheat, the nuclear photosynthetic ST-LS1 promoter from potato, the serine/threonine kinase (PAL) promoter and the glucoamylase (CHS) promoter from *Arabidopsis thaliana*. Also reported to be active in photosynthetically active tissues are the ribulose-1,5-bisphosphate carboxylase (RbcS) promoter from eastern larch (*Larix laricina*), the promoter for the *cab* gene, *cab6*, from pine, the promoter for the *Cab-1* gene from wheat, the promoter for the *CAB-1* gene from spinach, the promoter for the *cab1R* gene from rice, the pyruvate, orthophosphate dikinase (PPDK) promoter from corn, the promoter for the tobacco *Lhcb1*2* gene, the *Arabidopsis thaliana* SUC2 sucrose-H⁺ symporter and the promoter for the thylakoid membrane proteins from spinach (*psaD*, *psaF*, *psaE*, *PC*, *FNR*, *atpC*, *atpD*, *cab*, *rbcS*). Other promoters for the chlorophyll *a/b*-binding proteins may also be utilised in the invention, such as the promoters for *Lhcb* gene and *PsbP* gene from white mustard.

[0177] For the purpose of expression in sink tissues of the plant, such as the tuber of the potato plant, the fruit of tomato, or the seed of corn, wheat, rice and barley, it is desirable that the promoters driving expression of the target gene have relatively high expression in these specific tissues. A number of promoters for genes with tuber-specific or tuber-enhanced expression are known, including the class I patatin promoter, the promoter for the potato tuber ADPGPP genes, both the large and small subunits, the sucrose synthase promoter, the promoter for the major tuber proteins including the 22 kd protein complexes and protease inhibitors, the promoter for the granule-bound starch synthase gene (GBSS) and other class I and II patatins promoters.

[0178] Other promoters can also be used to express a target gene in specific tissues, such as seeds or fruits. Examples of such promoters include the 5' regulatory regions from such genes as napin, phaseolin, soybean trypsin inhibitor, ACP, stearyl-ACP desaturase, soybean α' subunit of β -conglycinin (soy 7s), and oleosin. Further examples include the promoter for β -conglycinin. Also included are the zeins, which are a group of storage proteins found in corn endosperm. Genomic clones for zein genes have been isolated and the promoters from these clones, including the 15 kD, 16 kD, 19 kD, 22 kD, 27 kD and genes, could also be used. Other promoters known to function, for example, in corn include the promoters for the following genes: waxy, Brittle, Shrunken 2, Branching enzymes I and II, starch synthases, debranching enzymes, oleosins, glutelins and sucrose synthases. Examples of promoters suitable for expression in wheat

include those promoters for the ADPglucose pyrosynthase (ADPGPP) subunits, the granule bound and other starch synthase, the branching and debranching enzymes, the embryogenesis-abundant proteins, the gliadins and the glutenins. Examples of such promoters in rice include those promoters for the ADPGPP subunits, the granule bound and other starch synthase, the branching enzymes, the debranching enzymes, sucrose synthases and the glutelins. Examples of such promoters for barley include those for the ADPGPP subunits, the granule bound and other starch synthase, the branching enzymes, the debranching enzymes, sucrose synthases, the hordeins, the embryo globulins and the aleurone specific proteins.

10 [0179] Root specific promoters may also be used. An example of such a promoter is the promoter for the acid chitinase gene. Expression in root tissue could also be accomplished using the root specific subdomains of the CaMV35S promoter that have been identified.

[0180] Desirable promoters for expression in cultured cells are strong constitutive promoters, or promoters that respond to a specific inducer (Gatz and Lenk, 1998, *Trends Plant Science* 3: 352-8). In certain embodiments, nucleic acid constructs expressing R polynucleotides of the present invention are introduced into banana plants that are susceptible Exemplary constitutive promoters for expression in intact banana plants are described in International Publication No. WO 02/053744 and in co-pending
20 PCT Application No. PCT/AU03/00919.

6.3.2 3' Non-translated region

[0181] The constructs of the present invention can comprise a 3' non-translated sequence. A 3' non-translated sequence refers to that portion of a gene comprising a DNA segment that contains a polyadenylation signal and any other regulatory signals capable of effecting mRNA processing or gene expression. The polyadenylation signal is characterised by effecting the addition of polyadenylic acid tracts to the 3' end of the mRNA precursor. Polyadenylation signals are commonly recognised by identity with the canonical form 5' AATAAA-3' although variations are not uncommon.

[0182] The 3' non-translated regulatory DNA sequence preferably includes from about 50 to 1,000 nucleotide base pairs and may contain plant transcriptional and translational termination sequences in addition to a polyadenylation signal and any other regulatory signals capable of effecting mRNA processing or gene expression. Examples

of suitable 3' non-translated sequences are the 3' transcribed non-translated regions containing a polyadenylation signal from the nopaline synthase (*nos*) gene of *Agrobacterium tumefaciens* (Bevan *et al.*, 1983, *Nucl. Acid Res.*, 11:369) and the terminator for the T7 transcript from the octopine synthase gene of *Agrobacterium tumefaciens*. Alternatively, suitable 3' non-translated sequences may be derived from plant genes such as the 3' end of the protease inhibitor I or II genes from potato or tomato, the soybean storage protein genes and the pea E9 small subunit of the ribulose-1,5-bisphosphate carboxylase (ssRUBISCO) gene, although other 3' elements known to those of skill in the art can also be employed. Alternatively, 3' non-translated regulatory sequences can be obtained *de novo* as, for example, described by An (1987, *Methods in Enzymology*, 153:292).

6.3.3 Optional sequences

[0183] The nucleic acid construct of the present invention can further include enhancers, either translation or transcription enhancers, as may be required. These enhancer regions are well known to persons skilled in the art, and can include the ATG initiation codon and adjacent sequences. The initiation codon must be in phase with the reading frame of the coding sequence relating to the foreign or endogenous DNA sequence to ensure translation of the entire sequence. The translation control signals and initiation codons can be of a variety of origins, both natural and synthetic. Translational initiation regions may be provided from the source of the transcriptional initiation region, or from the foreign or endogenous DNA sequence. The sequence can also be derived from the source of the promoter selected to drive transcription, and can be specifically modified so as to increase translation of the mRNA.

[0184] Examples of transcriptional enhancers include, but are not restricted to, elements from the CaMV 35S promoter and octopine synthase genes as for example described by Last *et al.* (U.S. Patent No. 5,290,924, which is incorporated herein by reference). It is proposed that the use of an enhancer element such as the *ocs* element, and particularly multiple copies of the element, will act to increase the level of transcription from adjacent promoters when applied in the context of plant transformation. Alternatively, the omega sequence derived from the coat protein gene of the tobacco mosaic virus (Gallie *et al.*, 1987) may be used to enhance translation of the mRNA transcribed from a polynucleotide according to the invention.

[0185] As the DNA sequence inserted between the transcription initiation site and the start of the coding sequence, i.e., the untranslated leader sequence, can influence gene expression, one can also employ a particular leader sequence. Preferred leader sequences include those that comprise sequences selected to direct optimum expression of the foreign or endogenous DNA sequence. For example, such leader sequences include a preferred consensus sequence which can increase or maintain mRNA stability and prevent inappropriate initiation of translation as for example described by Joshi (1987, *Nucl. Acid Res.*, 15:6643), which is incorporated herein by reference. However, other leader sequences, e.g., the leader sequence of RTBV, have a high degree of secondary structure that is expected to decrease mRNA stability and/or decrease translation of the mRNA. Thus, leader sequences (i) that do not have a high degree of secondary structure, (ii) that have a high degree of secondary structure where the secondary structure does not inhibit mRNA stability and/or decrease translation, or (iii) that are derived from genes that are highly expressed in plants, will be most preferred.

[0186] Regulatory elements such as the sucrose synthase intron as, for example, described by Vasil *et al.* (1989, *Plant Physiol.*, 91:5175), the Adh intron I as, for example, described by Callis *et al.* (1987, *Genes Develop.*, II), or the TMV omega element as, for example, described by Gallie *et al.* (1989, *The Plant Cell*, 1:301) can also be included where desired. Other such regulatory elements useful in the practice of the invention are known to those of skill in the art.

[0187] Additionally, targeting sequences may be employed to target R polypeptide to an intracellular compartment within plant cells or to the extracellular environment. For example, a DNA sequence encoding a transit or signal peptide sequence may be operably linked to a sequence encoding the R polypeptide or biologically active portion thereof such that, when translated, the transit or signal peptide can transport the polypeptide or portion to a particular intracellular or extracellular destination, and can then be post-translationally removed. Transit or signal peptides act by facilitating the transport of proteins through intracellular membranes, e.g., endoplasmic reticulum, vacuole, vesicle, plastid, mitochondrial and plasmalemma membranes. For example, the targeting sequence can direct a desired protein to a particular organelle such as a vacuole or a plastid (e.g., a chloroplast), rather than to the cytosol. Thus, the nucleic acid construct can further comprise a plastid transit peptide encoding DNA sequence operably linked between a promoter region or promoter variant according to the

invention and the foreign or endogenous DNA sequence. For example, reference may be made to Heijne *et al.* (1989, *Eur. J. Biochem.*, 180:535) and Keegstra *et al.* (1989, *Ann. Rev. Plant Physiol. Plant Mol. Biol.*, 40:471), which are incorporated herein by reference.

- 5 [0188] The nucleic acid construct is typically introduced into a vector, such as a plasmid. Plasmid vectors include additional DNA sequences that provide for easy selection, amplification, and transformation of the expression cassette in prokaryotic and eukaryotic cells, *e.g.*, pUC-derived vectors, pSK-derived vectors, pGEM-derived vectors, pSP-derived vectors, or pBS-derived vectors. Additional DNA sequences
- 10 include origins of replication to provide for autonomous replication of the vector, selectable marker genes, preferably encoding antibiotic or herbicide resistance, unique multiple cloning sites providing for multiple sites to insert DNA sequences or genes encoded in the nucleic acid construct, and sequences that enhance transformation of prokaryotic and eukaryotic cells.
- 15 [0189] The vector desirable contains an element(s) that permits either stable integration of the vector into the host cell genome or autonomous replication of the vector in the cell independent of the genome of the cell. The vector may be integrated into the host cell genome when introduced into a host cell. For integration, the vector may rely on a foreign or endogenous DNA sequence present therein or any other
- 20 element of the vector for stable integration of the vector into the genome by homologous recombination. Alternatively, the vector may contain additional nucleic acid sequences for directing integration by homologous recombination into the genome of the host cell. The additional nucleic acid sequences enable the vector to be integrated into the host cell genome at a precise location in the chromosome. To increase the
- 25 likelihood of integration at a precise location, the integrational elements should preferably contain a sufficient number of nucleic acids, such as 100 to 1,500 base pairs, preferably 400 to 1,500 base pairs, and most preferably 800 to 1,500 base pairs, which are highly homologous with the corresponding target sequence to enhance the probability of homologous recombination. The integrational elements may be any
- 30 sequence that is homologous with the target sequence in the genome of the host cell. Furthermore, the integrational elements may be non-encoding or encoding nucleic acid sequences.

[0190] For cloning and subcloning purposes, the vector may further comprise an origin of replication enabling the vector to replicate autonomously in a host cell such as a bacterial cell. Examples of bacterial origins of replication are the origins of replication of plasmids pBR322, pUC19, pACYC177, and pACYC184 permitting replication in *E. coli*, and pUB110, pE194, pTA1060, and pAM β 1 permitting replication in *Bacillus*. The origin of replication may be one having a mutation to make its function temperature-sensitive in a *Bacillus* cell (see, e.g., Ehrlich, 1978, *Proc. Natl. Acad. Sci. USA* 75:1433).

6.3.4 Marker genes

10 [0191] To facilitate identification of transformants, the nucleic acid construct desirably comprises a selectable or screenable marker gene as, or in addition to, a polynucleotide sequence according to the invention. The actual choice of a marker is not crucial as long as it is functional (i.e., selective) in combination with the plant cells of choice. The marker gene and the R polynucleotide sequence of interest do not have to
15 be linked, since co-transformation of unlinked genes as, for example, described in U.S. Pat. No. 4,399,216 is also an efficient process in plant transformation.

[0192] Included within the terms selectable or screenable marker genes are genes that encode a "secretable marker" whose secretion can be detected as a means of identifying or selecting for transformed cells. Examples include markers that encode a
20 secretable antigen that can be identified by antibody interaction, or secretable enzymes that can be detected by their catalytic activity. Secretable proteins include, but are not restricted to, proteins that are inserted or trapped in the cell wall (e.g., proteins that include a leader sequence such as that found in the expression unit of extensin or tobacco PR-S); small, diffusible proteins detectable, e.g. by ELISA; and small active
25 enzymes detectable in extracellular solution (e.g., α -amylase, β -lactamase, phosphinothricin acetyltransferase).

6.3.5 Selectable markers

[0193] Examples of bacterial selectable markers are the *dal* genes from *Bacillus subtilis* or *Bacillus licheniformis*, or markers that confer antibiotic resistance such as
30 ampicillin, kanamycin, erythromycin, chloramphenicol or tetracycline resistance. Exemplary selectable markers for selection of plant transformants include, but are not limited to, a *hyg* gene which encodes hygromycin B resistance; a neomycin

phosphotransferase (*neo*) gene conferring resistance to kanamycin, paromomycin, G418 and the like as, for example, described by Potrykus *et al.* (1985, *Mol. Gen. Genet.* 199:183); a glutathione-S-transferase gene from rat liver conferring resistance to glutathione derived herbicides as, for example, described in EP-A 256 223; a glutamine synthetase gene conferring, upon overexpression, resistance to glutamine synthetase inhibitors such as phosphinothricin as, for example, described WO87/05327, an acetyl transferase gene from *Streptomyces viridochromogenes* conferring resistance to the selective agent phosphinothricin as, for example, described in EP-A 275 957, a gene encoding a 5-enolshikimate-3-phosphate synthase (EPSPS) conferring tolerance to N-phosphonomethylglycine as, for example, described by Hinchee *et al.* (1988, *Biotech.*, 6:915), a *bar* gene conferring resistance against bialaphos as, for example, described in WO91/02071; a nitrilase gene such as *bxn* from *Klebsiella ozaenae* which confers resistance to bromoxynil (Stalker *et al.*, 1988, *Science*, 242:419); a dihydrofolate reductase (DHFR) gene conferring resistance to methotrexate (Thillet *et al.*, 1988, *J. Biol. Chem.*, 263:12500); a mutant acetolactate synthase gene (ALS), which confers resistance to imidazolinone, sulfonylurea or other ALS-inhibiting chemicals (EP-A-154 204); a mutated anthranilate synthase gene that confers resistance to 5-methyl tryptophan; or a dalapon dehalogenase gene that confers resistance to the herbicide.

6.3.6 Screenable markers

[0194] Preferred screenable markers include, but are not limited to, a *uidA* gene encoding a β -glucuronidase (GUS) enzyme for which various chromogenic substrates are known; a β -galactosidase gene encoding an enzyme for which chromogenic substrates are known; an aequorin gene (Prasher *et al.*, 1985, *Biochem. Biophys. Res. Comm.*, 126:1259), which may be employed in calcium-sensitive bioluminescence detection; a green fluorescent protein gene (Niedz *et al.*, 1995 *Plant Cell Reports*, 14:403); a luciferase (*luc*) gene (Ow *et al.*, 1986, *Science*, 234:856), which allows for bioluminescence detection; a β -lactamase gene (Sutcliffe, 1978, *Proc. Natl. Acad. Sci. USA* 75:3737), which encodes an enzyme for which various chromogenic substrates are known (e.g., PADAC, a chromogenic cephalosporin); an R-locus gene, encoding a product that regulates the production of anthocyanin pigments (red colour) in plant tissues (Dellaporta *et al.*, 1988, in *Chromosome Structure and Function*, pp. 263-282); an α -amylase gene (Ikuta *et al.*, 1990, *Biotech.*, 8:241); a tyrosinase gene (Katz *et al.*, 1983, *J. Gen. Microbiol.*, 129:2703) which encodes an enzyme capable of oxidising

tyrosine to dopa and dopaquinone which in turn condenses to form the easily detectable compound melanin; or a *xylE* gene (Zukowsky *et al.*, 1983, *Proc. Natl. Acad. Sci. USA* 80:1101), which encodes a catechol dioxygenase that can convert chromogenic catechols.

5 **7. Introduction of the nucleic acid construct into plant cells**

[0195] The sequences of the present invention can be used to transform or transfect any plant. In this manner, genetically modified plants, plant cells, plant tissue, seed, and the like can be obtained. Transformation protocols as well as protocols for introducing nucleotide sequences into plants may vary depending on the type of plant or plant cell,
10 i.e. monocot or dicot, targeted for transformation. It is recognised that the transformation protocols may be used for transfection or introduction of the oligonucleotide sequences to disrupt R gene function. Suitable methods of introducing nucleotide sequences into plant cells and subsequent insertion into the plant genome include microinjection (Crossway *et al.*, 1986, *Biotechniques* 4:320-334),
15 electroporation (Riggs *et al.*, 1986, *Proc. Natl. Acad. Sci. USA* 83:5602-5606), Agrobacterium-mediated transformation (Townsend *et al.*, U.S. Pat. No. 5,563,055; Zhao *et al.*, U.S. Pat. No. 5,981,840), direct gene transfer (Paszkowski *et al.*, 1984, *EMBO J.* 3:2717-2722), and ballistic particle acceleration (see, for example, Sanford *et al.*, U.S. Pat. No. 4,945,050; Tomes *et al.*, U.S. Pat. No. 5,879,918; Tomes *et al.*, U.S.
20 Pat. No. 5,886,244; Bidney *et al.*, U.S. Pat. No. 5,932,782; Tomes *et al.* (1995) "Direct DNA Transfer into Intact Plant Cells via Microprojectile Bombardment," in *Plant Cell, Tissue, and Organ Culture: Fundamental Methods*, ed. Gamborg and Phillips (Springer-Verlag, Berlin); and McCabe *et al.*, (1988, *Biotechnology* 6:923-926). Also see Weissinger *et al.* (1988 *Ann. Rev. Genet.* 22:421-477), Sanford *et al.*, (1987, *Particulate*
25 *Science and Technology* 5:27-37; onion), Christou *et al.*, (1988, *Plant Physiol.* 87:671-674; soybean); Datta *et al.*, (1990, *Biotechnology* 8:736-740; rice), Klein *et al.* (1988, *Proc. Natl. Acad. Sci. USA* 85:4305-4309, maize), Hooykaas-Van Slogteren *et al.* (1984, *Nature (London)* 311:763-764; cereals), Bowen *et al.*, (U.S. Pat. No. 5,736,369; cereals), Bytebier *et al.*, (1987, *Proc. Natl. Acad. Sci. USA* 84:5345-5349; Liliaceae),
30 De Wet *et al.* (1985, in *The Experimental Manipulation of Ovule Tissues*, ed. Chapman *et al.* (Longman, N.Y.), pp. 197-209; pollen), Kaeppler *et al.*, (1990, *Plant Cell Reports* 9:415-418; 1992, *Theor. Appl. Genet.* 84:560-566; whisker-mediated transformation), D'Halluin *et al.* (1992, *Plant Cell* 4:1495-1505; electroporation); Li *et al.*, (1993, *Plant*

Cell Reports 12:250-255; rice), Christou and Ford (1995, *Annals of Botany* 75:407-413; rice) and Osjoda *et al.* (1996, *Nature Biotechnology* 14:745-750; maize via *Agrobacterium tumefaciens*). Guidance in the practical implementation of transformation systems for plant improvement is provided by Birch (1997, *Annu. Rev. Plant Physiol. Plant Molec. Biol.* 48: 297-326).

[0196] In certain embodiments, the present invention is concerned with transforming monocotyledonous plants, including graminaceous and non-graminaceous monocotyledonous plants. Illustrative examples of non-graminaceous monocotyledonous plants include, but are not limited to, *Musaceae* (*Musa* and *Ensete*), taro, ginger, onions, garlic, pineapple, bromeliads, palms, orchids, lilies, irises and the like. There are a variety of methods known currently for transformation of monocotyledonous plants. Presently, preferred methods for transformation of monocots are microprojectile bombardment of explants or suspension cells, and direct DNA uptake or electroporation as, for example, described by Shimamoto *et al.* (1989, *supra*).

Transgenic maize plants have been obtained by introducing the *Streptomyces hygroscopicus bar* gene into embryogenic cells of a maize suspension culture by microprojectile bombardment (Gordon-Kamm, 1990, *Plant Cell*, 2:603-618). The introduction of genetic material into aleurone protoplasts of other monocotyledonous crops such as wheat and barley has been reported (Lee, 1989, *Plant Mol. Biol.* 13:21-30). Wheat plants have been regenerated from embryogenic suspension culture by selecting only the aged compact and nodular embryogenic callus tissues for the establishment of the embryogenic suspension cultures (Vasil, 1990, *Bio/Technol.* 8:429-434). The combination with transformation systems for these crops enables the application of the present invention to monocots. These methods may also be applied for the transformation and regeneration of dicots. Transgenic sugarcane plants have been regenerated from embryogenic callus as, for example, described by Bower *et al.* (1996, *Molecular Breeding* 2:239-249).

8. Production and characterisation of differentiated transgenic plants

8.1 Regeneration

[0197] The methods used to regenerate transformed cells into differentiated plants are not critical to this invention, and any method suitable for a target plant can be employed. Normally, a plant cell is regenerated to obtain a whole plant following a transformation process.

- [0198] Regeneration from protoplasts varies from species to species of plants, but generally a suspension of protoplasts is made first. In certain species, embryo formation can then be induced from the protoplast suspension, to the stage of ripening and germination as natural embryos. The culture media will generally contain various amino acids and hormones, necessary for growth and regeneration. Examples of hormones utilised include auxins and cytokinins. It is sometimes advantageous to add glutamic acid and proline to the medium, especially for such species as corn and alfalfa. Efficient regeneration will depend on the medium, on the genotype, and on the history of the culture. If these variables are controlled, regeneration is reproducible. Regeneration also occurs from plant callus, explants, organs or parts. Transformation can be performed in the context of organ or plant part regeneration as, for example, described in *Methods in Enzymology*, Vol. 118 and Klee *et al.* (1987, *Annual Review of Plant Physiology*, 38:467), which are incorporated herein by reference. Utilising the leaf disk-transformation-regeneration method of Horsch *et al.* (1985, *Science*, 227:1229, incorporated herein by reference), disks are cultured on selective media, followed by shoot formation in about 2-4 weeks. Shoots that develop are excised from calli and transplanted to appropriate root-inducing selective medium. Rooted plantlets are transplanted to soil as soon as possible after roots appear. The plantlets can be repotted as required, until reaching maturity.
- 20 [0199] In vegetatively propagated crops, the mature transgenic plants are propagated by the taking of cuttings or by tissue culture techniques to produce multiple identical plants. Selection of desirable transgenes is made and new varieties are obtained and propagated vegetatively for commercial use.
- 25 [0200] In seed propagated crops, the mature transgenic plants can be self-crossed to produce a homozygous inbred plant. The inbred plant produces seed containing the newly introduced foreign gene(s). These seeds can be grown to produce plants that would produce the selected phenotype, e.g., early flowering.
- 30 [0201] Parts obtained from the regenerated plant, such as flowers, seeds, leaves, branches, fruit, and the like are included in the invention, provided that these parts comprise cells that have been transformed as described. Progeny and variants, and mutants of the regenerated plants are also included within the scope of the invention, provided that these parts comprise the introduced nucleic acid sequences.

[0202] The literature describes numerous techniques for regenerating specific plant types and more are continually becoming known. Those of ordinary skill in the art can refer to the literature for details and select suitable techniques without undue experimentation.

5 8.2 Characterisation

[0203] To confirm the presence of a R polynucleotide of the invention in the regenerating plants, a variety of assays may be performed. Such assays include, for example, "molecular biological" assays well known to those of skill in the art, such as Southern and Northern blotting and PCR; an R protein expressed by the polynucleotide
10 of the invention may be assayed using antigen-binding molecules as for example described herein.

[0204] In order that the invention may be readily understood and put into practical effect, particular preferred embodiments will now be described by way of the following non-limiting example.

EXAMPLES

EXAMPLE 1

Identification of R genes from *M. acuminata*

CTAB total DNA purification

5 [0205] Genomic DNA was extracted using the CTAB protocol of Stewart & Via (1993). Briefly, leaf tissue was frozen in liquid N₂ and ground in a mortar and pestle. Powdered tissue was resuspended in CTAB Buffer (1% Sarcosine, 0.8 M NaCl, 0.022 M EDTA pH8.0, 0.22 M Tris-HCl pH 7.8, 0.8% CTAB, 0.14 M Mannitol) at 65°C. An equal volume of chloroform:isoamylalcohol (24:1) was immediately added, mixed by
10 inversion and incubated at 65°C for 10 min with occasional inversion. Samples were centrifuged for 5 min at 14000 rpm in a microfuge to separate phases. The aqueous layer was collected and an equal volume of isopropanol added. DNA was spooled out, washed in 70% ethanol, and allowed to dry before resuspending in 100 µL dH₂O containing RNaseA (1 mg/mL).

15 *Purification of total RNA*

[0206] Total RNA extractions were performed using the method of Chang *et al.* (1993). Tissue was frozen in liquid N₂ and ground to a powder in a mortar and pestle. Powdered tissue was added to preheated (65° C) extraction buffer (2% CTAB, 2% PVP, 100 mM Tris HCl pH8, 25 mM EDTA, 2 M NaCl, 0.05% spermidine, 2% beta-
20 mercaptoethanol). Chloroform:isoamylalcohol (24:1) was added, the suspension vortexed, and samples centrifuged at top speed in a microfuge for 5 min. The aqueous phase was collected and an equal volume of DEPC-treated 4M LiCl added. RNA was precipitated overnight 4° C and then centrifuged at 4° C for 30 min at top speed. Pelleted RNA was resuspended in 10 X SSTE and extracted once more with
25 chloroform:isoamylalcohol (24:1). The RNA was reprecipitated at -20° C overnight following the addition of 1/10 volume DEPC-treated 2.5 M NaOAc pH6.0 and 21/2 volumes of 100% ethanol. Tubes were centrifuged 20 min, the pellets washed with 70% ethanol and resuspended in DEPC-treated dH₂O.

Reverse-transcriptase PCR of banana R-genes

30 [0207] Sequences of R-genes from plant species were aligned and degenerate primers designed to conserved motifs in the NBS regions. The degenerate primers were

used to generate single-stranded cDNAs from total RNA using reverse transcriptase and then to subsequently amplify the NBS region of the banana R-genes. To generate the region 5' of the NBS domain, RNA primers were ligated to the 5' end of the mRNA after removal of the 5'-cap structure. Ligated mRNA was reverse transcribed using reverse transcriptase to generate single-stranded cDNA. Primer complementary to the ligated RNA primer and a specific primer to the known NBS sequence was added and PCR undertaken to generate the 5' region of the R-gene using the parameters of: initial denaturation step of 94° C for 2 min followed by 5 cycles of 94° C for 30 secs, 55-65° C for 30 secs, 72° C for 3-5 min, then 25 cycles of 94° C for 30 secs, 45-60° C for 30 secs, 72° C for 3-5 min, followed by a final annealing step 72° C for 10 min. N-terminal and C-terminal primers were subsequently used to amplify complete R-gene sequences from genomic DNA using PCR with the following conditions: initial denaturation step of 94° C for 2 min, followed by 25 cycles of 94° C for 30 secs, 55° C for 30 secs, 72° C for 1-5 min, followed by a final annealing step 72° C for 10 min. All PCR products were cloned and sequenced to verify identity. The full-length nucleotide sequences for two R genes, one isolated from *Musa acuminata* (Calcutta 4) designated RGA5 and the other from *Musa acuminata* spp *malaccensis* designated RGA2, are presented in SEQ ID NO: 1 and 3, respectively.

[0208] RT-PCR was then used to compare the expression of the R genes between *M. acuminata* spp *malaccensis* plants that were susceptible or resistant to *Fusarium oxysporum* fsp *cubense* (FOC). The results presented in Figure 3 show that the RGA2 gene (see lanes C2) is transcribed in FOC resistant plants but not in FOC sensitive plants. This suggests that RGA2 may be an attractive candidate for conferring disease resistance to susceptible plants. The inventors propose to transform Cavendish, which is resistant to race 1 but susceptible to race 4, (i) with RGA2 only; (ii) with RGA5 and (iii) with both RGA2 and RGA5, under the control of a heterologous promoter (e.g., Ubi) or the native RGA2 promoter.

The disclosure of every patent, patent application, and publication cited herein is hereby incorporated herein by reference in its entirety.

[0209] The citation of any reference herein should not be construed as an admission that such reference is available as "Prior Art" to the instant application.

5 [0210] Those skilled in the art will appreciate that the invention described herein is susceptible to variations and modifications other than those specifically described. It is to be understood that the invention includes all such variations and modifications. The invention also includes all of the steps, features, compositions and compounds referred to or indicated in this specification, individually or collectively, and any and all
10 combinations of any two or more of said steps or features.

DATED this 25 September, 2003

QUEENSLAND UNIVERSITY OF TECHNOLOGY

By their Patent Attorneys

DAVIES COLLISON CAVE

15

BIBLIOGRAPHY

- Aarts N, Metz M, Holub E, Staskawic B, Daniels M and Parker J (1998).** Different requirements for EDS1 and NDR1 by disease resistance genes define at least two R gene-mediated signalling pathways in Arabidopsis. *Proc. Natl. Acad. Sci. USA.* 95, 10306-10311.
- Aarts M, Lintel G, Hekkert B, Holub E, Beynon J, Stiekema W and Pereira A (1998).** Identification of R-gene homologous DNA fragments genetically linked to disease resistance loci in Arabidopsis thaliana. *Mol. Plant-Microbe Interact.* 11: 251-258.
- Anderson P, Lawrence G, Morrish B, Finnegan E and Ellis J (1997).** Inactivation of the flax rust resistance gene M associated with loss of a repeated unit within the leucine-rich repeat coding region. *Plant cell.* 9: 641-651.
- Aravind L, Dixit V and Koonin E (1999).** The domains of the death: evolution of the apoptosis machinery. *Trends in biochemical science.* 24: 47-53.
- Asai T, Tena G, Plotnikova J, Willmann M, Chiu W, Gomez-Gomez L, Boller T, Ausubel M and Sheen J (2002).** MAP kinase signalling cascade in Arabidopsis innate immunity. *Nature.* 415, 977-983.
- Baker B, Zambryski P, Staskawicz B and Dinesh-Kumar S (1997).** Signaling in plant-microbe interactions. *Science.* 276, 726-733.
- Ballvora A, Ercolano M, Weib J, Meksem K, Bormann C, Oberhagemann P, Salamini F and Gebhardt C (2002).** The R1 gene for potato resistance to late blight (*Phytophthora infestans*) belongs to the leucine zipper/NBS/LRR class of plant resistance gene. *The Plant Journal.* 30, 361-371.
- Becker D, Dugdale B, Smith M, Harding R and Dale J (2000).** Genetic transformation of Cavendish banana (*Musa* spp. AAA group) cv 'Grand Nain' via microprojectile bombardment. *Plant Cell Reports.* 19, 229-234.
- Bendahmane A, Kanyuka K and Baulcombe D (1999).** The Rx gene from tomato controls separate virus resistance and cell death responses. *Plant Cell.* 11: 781-791.

- Bentley S, Pegg K, Moore N, Davis R and Buddenhagen W (1998).** Genetic variation among vegetative compatibility groups of *Fusarium oxysporum* f. sp. *cubense* analysed by DNA fingerprinting . *Phytopathology* 88: 1283-1293.
- Bittner-Eddy P, Crute I, Holub E and Beynon J (2000).** RPP13 is a simple locus in *Arabidopsis thaliana* for alleles that specify downy mildew resistance to different avirulence determinants in *peronospora parasitica*. *The Plant Journal*. 21, 177-188.
- Bonas U and Lahaye T (2002).** Plant disease resistance triggered by pathogen-derived molecules: refined models of specific recognition. *Current opinion in microbiology*. 5: 44-50.
- Botella M, Pareker J, Frost L, Bittner-Eddy, Beynon J et al. (1998).** Three genes of the *Arabidopsis* RPP1 complex resistance locus recognize distinct *Peronospora parasitica* avirulence determinants. *Plant cell* 10: 1847-1860.
- Brommonschenkel S, Fray A and Tanksley S (2000).** The broad-spectrum tospovirus resistance gene Sw-5 of tomato is a homolog of the root-knot nematode resistance gene Mi. *Molecular Plant Microbe Interactioin*. 13, 1130-1138.
- Brueggeman R, Rostoks N, Kudrna D, Killian A, Han F, Chen J, Druka A, Steffenson B and Kleinhofs A (2002).** The barley steam rust-resistance gene Rpg1 is a novel disease-resistance gene with homology to receptor kinases. *PNAS*. 1-6.
- Bryan G, Wu K, Farall L, Jia Y, Hershey H, McAdams S, Faulk K, Donaldson G, Tarchini R and Valent B (2000).** A single amino acid difference distinguishes resistant and susceptible alleles of the rice blast resistance gene Pi-ta. *Plant Cell*. 12: 2033-2045.
- Cai D, Kleine M, Kiffle S, Harloff H, Sanda N, Marcker K, Klein-Lankhorts R, Salentjn E, Lange W, Stiekema W, Wyss W, Grundler F and Jung C (1997).** Positional cloning of a gene for nematode resistance in sugar beet. *Science*. 275, 832-834.
- Cao H, Glazebrook J, Clarke J, Volko S and Dong X (1997).** The *Arabidopsis* NPR1 gene that controls systemic acquired resistance encodes a novel protein containing ankyrin repeats. *Cell*. 88, 57-63.

FAO,

2001<http://apps.fao.org/lim500/nph-wrap.pi?production.crops.primary&Domain=SU&>

- Cao H, Li X and Dong X (1998).** Generation of broad-spectrum disease resistance by overexpression of an essential regulator gene in systemic acquired resistance. *Proc Natl Acad Sci USA*. 95, 6531-6536.
- Century K, Shapiro A, Repetti P, Dahlbeck D, Holub E and Staskawicz J (1997).** NDR1, a pathogen-induced component required for Arabidopsis disease resistance. *Science*. 278: 1963-1965.
- Chern M, Fitzgerald H, Yadav R, Canlas P, Dong X and Ronald P (2001).** Evidence for a disease-resistance pathway in rice similar to the NPR1-mediated signalling pathway in Arabidopsis. *The plant Journal*. 27, 101-113.
- Cohn J, Sessa G and Martin G (2001).** Innate immunity in plants. *Current Opinoin in Immunology*. 13, 55-62.
- Collins N, Drake J, Ayliffe M, Sun Q, Ellis J, Hulbert S, and Pryor T (1999).** Molecular characterization of the Maize Rpl-D Rust resistance haplotype and its mutants. *The Plant Cell*. 11, 1365-1376.
- Despres C, DeLong C, Glaze S, Liu E and Fobert P (2000).** The Arabidopsis NPR1/NIM1 protein enhances the DNA binding activity of a subgroup of the TGA family of bZIP transcription factors. *The Plant Cell*. 12, 279-290.
- Dixon M, Hatzixanthis K, Jones D, Harrison K, Jones J (1998).** The tomato Cf-5 disease resistance gene and six homologs show pronounced allelic variation in leucine-rich repeat copy number. *Plant Cell*. 10, 1915-1925.
- Dodds P, Lawrence G and Ellis J (2000).** Six amino acid changes confined to the Leucine-Rich repeat B-strand/B-turn motif determine the difference between the P and P2 rust resistance specificities in flax. *Plant Cell* 13: 163-178.
- Ellis J, Dodds P and Pryor T (2000).** The generation of plant disease resistance gene specificities. *Trends in Plant Science*. 5, 373-379.
- Ellis J and Jones D (1998).** Structure and function of proteins controlling strain-specific pathogen resistance in plants. *Current opinion in plant biology*. 1, 288-293.

- Endo T, Ikeo K and Gojobori T (1996).** Large-scale search for genes on which positive selection may operate. *Molecular Biology Evolution*. 13, 685-690.
- Ernst K, Kumar A, Kriseleit D, Kloos D, Phillips M and Ganai M (2002).** The broad-spectrum potato cyst nematode resistance gene (Hcr) from tomato is the only member of a large gene family of NBS-LRR genes with an unusual amino acid repeat in the LRR region. *Plant Journal*. 31: 127-136.
- Falk A, Feys B, Frost L, Jones J, Daniels M and Parker J (1999).** EDS1, an essential component of R gene-mediated disease resistance in *Arabidopsis* has homology to eukaryotic lipases. *Proc. Natl. Acad. Sci. USA*. 96: 3292-3297.
- Glazebrook J (2001).** Genes controlling expression of defence responses in *Arabidopsis*. 4, 301-308.
- Graham M, Mareck L, Lohnes D, Cregan P and Shoemaker R (2000).** Expression and genome organization of resistance gene analogues in soybean. *Genome*. 43, 86-93.
- Grant M, Godard L, Straube E, Ashfield T, Leward J, Sattler A, Innes R and Dangl J (1995).** Structure of the *Arabidopsis* RPM1 enabling dual specificity disease resistance. *Science*. 269, 843-846.
- Gassmann W, Hinsch M and Staskawicz B (1999).** The *Arabidopsis* RPS4 bacterial-resistance gene is a member of the TIR-NBS-LRR family of disease resistance genes. *Plant Journal*. 20: 265-277.
- Gomez-Gomez L and Boller T (2000).** FLS2: An LRR receptor-like kinase involved in the perception of the bacterial elicitor flagellin in *Arabidopsis*. *Mol. Cell*. 5, 1003-1011.
- Hammond-Kosack K and Jones J (1997).** Plant resistance genes. *Annu. Rev. Plant. Physiol. Plant Mol. Biol.* 48, 575-607.
- Hughes L and Nei M (1988).** Pattern of nucleotide substitution at major histocompatibility complex class I loci reveals overdominant selection. *Nature*. 335, 167-170.
- Hulbert S, Webb C, Smith S and Sun Q (2001).** Resistance gene complexes: evolution and utilization. *Annu. Rev. Phytopathol.* 39: 285-312.

- Jia Y, McAdams S, Bryan G, Hershey P and Valent B (2000).** Direct interaction of resistance gene and avirulence gene products confers rice blast resistance. *EMBO J.* 19, 4004-4014.
- Johal G, and Briggs S (1992).** Reductase activity encoded by the HM1 disease resistance gene in maize. *Science.* 258, 985-987.
- Kanazin V, Mareck L and Shoemaker R (1996).** Resistance gene analogs are conserved and clustered in soybean. *Proc Natl Acad Sci USA.* 93, 11746-11750.
- Kawchuk L, Hachey J, Lynch D, Kulcsar F, Rooijen G, Waterer D, Robertson A, Kokko E, Byers R, Howard R, Fischer R and Prufer D (2001).** Tomato Ve disease resistance genes encode cell surface-like receptors. *PNAS.* 98, 6511-6515.
- Kimura M (1983).** The neutral theory of molecular evolution. Cambridge University Press.
- Kinkema M, Fan W and Dong X (2000).** Nuclear localization of NPR1 is required for activation of PR gene expression. *The plant cell.* 12, 2339-2350.
- Lagudah E, Moullet O and Appels R (1997).** Map-based cloning of a resistance gene sequence encoding a nucleotide-binding domain and leucine rich region at the Cre3 nematode resistance locus of wheat. *Genome.* 40, 659-665.
- Lawrence GJ, Finnegan EJ, Ayliffe M and Ellis J (1995).** The L6 gene for flax rust resistance is related to the Arabidopsis bacterial resistance gene RPS2 and the tobacco viral resistance gene N. *Plant Cell.* 7: 1195-1206.
- Leister D, Ballvora A, Salamini F and Gebhardt C (1996).** A PCR-based approach for isolating pathogen resistance genes from potato with potential for wide application in plants. *Nature Genetics:* 14, 421-429.
- Leister D, Kurth J, Laurie D, Yano M, Sasaki T, Devos K, Graner A and Schulze-Lefert P (1998).** Rapid organization of resistance gene homologues in cereal genomes. *Proc Natl Acad Sci USA.* 95, 370-375.
- May G, Afza R, Mason H, Wiecko A, Novak F and Arntzen C (1995).** Generation of transgenic banana (*Musa acuminata*) plants via *Agrobacterium*-mediated transformation. *Bio/Technology.* 13, 486-492.

- Mes J, Van Doorn A, Wijbrandi J, Simons G, Cornelissen B and Haring M (2000).** Expression of the Fusarium resistance gene I-2 colocalizes with the site of fungal containment. *The plant journal*. 23, 183-193.
- Meyers B, Shen K, Rohani P, Gaut B and Michelmore R (1998).** Receptor -like genes in the mayor resistance locus in lettuce are subject of divergent selection. *Plant Cell*. 11, 1833-1846.
- Meyers B, Dickerman A, Michelmore R, Sivaramakrishnan S, Sobral B and Young N (1999).** Plant disease resistance genes encode members of an ancient and diverse protein family within the nucleotide-binding superfamily. *The Plant Journal*. 20, 317-332.
- Milligan S, Bodeau J, Yaghoobi J, Kaloshian I, Zabel P and Williamson V (1998).** The root knot resistance gene Mi from tomato is a member of the leucine zipper, nucleotide binding site, leucine-rich repeat family of plant genes. *The plant cell*. 10, 1307-1319.
- Mindrinis M, Katagiri F, Yu G and Ausubel F (1994).** The *A. thaliana* disease resistance gene RPS2 encodes a protein containing a nucleotide-binding site and leucine-rich repeats. *Cell*. 78: 1089-1099.
- Muskett P, Kahn K, Austin M, Moisan L, Sadanandom A, Shirasu K, Jones J and Parker J (2002).** Arabidopsis RAR1 exerts rate-limiting control of R genes-mediated defenses against multiple pathogens. *The plant cell*. 14. 979-992.
- Ortiz R, Ferris R and Vuylsteke D (1995).** Banana and plantain breeding. In banana and plantains. Gowen, S. Ed. Chapman and Hall London. pp, 110-146.
- Pan Q, Wendel J and Fluhr R (2000).** Divergent evolution of plant NBS-LRR resistance gene homologues in dicot and cereal genomes. *Journal of molecular evolution*. 50: 203-213.
- Parker J, Coleman M, Szabo V, Frost V, Schmidt R, Van der Biezen E, Moores T, Dean C, Daniels M and Jones J (1997).** The Arabidopsis downy mildew resistance gene RPP5 shares similarity to the toll interleukine-1 receptors with N and L6. *The Plant Cell*. 9, 879-894.
- Parniske M, Hammond-Kosack K, Golstein C, Thomas C, Jones D, Harrison K, Wulff BB, Jones J (1997).** Novel disease resistance specificities result from

- sequence exchange between tandemly repeated genes at the Cf-4/9 locus of tomato. *Cell*. 91, 821-832
- Ploetz R (2000).** Panama disease: a classical and destructive disease of banana. Online. Plant Health Progress doi: 10.1094/PHP-2000-1240-01-HM.
- Ploetz R and Pegg K (2000).** Fungal disease of the root, corm and pseudostem. In Diseases of banana, abaca and ensete. Jones, D. Ed. CABI.
- Richter T and Ronald P (2000).** The evolution of disease resistance genes. *Plant Molecular Biology*. 42, 195-204.
- Rommens C and Kishore G (2000).** Exploiting the full potential of disease-resistance genes for agricultural use. *Current opinion in biotechnology*. 11, 120-125.
- Ryals J, Weymann K, Lawton K, Friedrich L, Ellis D, Steiner H, Johnson J, Delaney T, Jesse T, Vos P and Uknes S (1997).** The Arabidopsis NIM1 protein shows homology to the mammalian transcription factor inhibitor I κ B. *The Plant Cell*. 9, 425-439.
- Sagi L, Panis B, Remy S, Schoofs H, De Smet K, Swennen R and Cammue P (1995).** Genetic transformation of banana and plantain (*Musa* spp) via particle bombardment. *Bio/Techniques*. 13, 481-485.
- Salmeron J, Oldroyd E, Rommens C, Scofield S, Kim H, Lavelle D, Dahlbeck D and Staskawicz B (1996).** Tomato Prf is a member of a leucine-rich repeat class of plant disease resistance gene and lies embedded within the Pto kinase gene cluster. *Cell*. 86, 123-133.
- Shen K, Meyers B, Islam-Faridi M, Chin D, Stelly D and Michelmore R (1998).** Resistance gene candidates identified by PCR with degenerate oligonucleotide primers Map to clusters of resistance genes in lettuce. *Molecular Plant Pathogen Interaction*. 11, 815-823.
- Shirasu K, Lahaye T, Tan M, Zhou F, Azevedo C and Schulze-Lefert P (1999).** A novel class of eukaryotic zinc-binding proteins is required for disease resistance signalling in barley and the development in *C. elegans*. *Cell*. 99: 355-366.
- Simons G, Groenendijk J, Wijbrandi J, Reijans M, Groenen J, Diergaarde P, Van der Lee T, Bleeker M, Onstenk J, Both M, Haring M, Mes J, Cornelisse B, Zabeau M and Vos P (1998).** Dissection of the Fusarium I2 gene cluster in

- tomato reveals six homologs and one active gene copy. *The Plant Cell*. 10, 1055-1068.
- Song W, Wang G, Gardner J, Holsten T, Ronald P (1997). Evolution of the rice Xa21 disease resistance gene family. *Plant Cell* 9: 1279-1287.
- Song W, Wang G, Chen L et al. (1998). A receptor kinase-like protein encoded by the rice disease Xa21. *Science*. 270, 1804-1806.
- Stuiver M and Custers J (2002). Engineering disease resistance in plants. *Nature*. 411: 865-868.
- Tai T, Dahlbeck D, Clark E, Gajiwala P, Pasion R, Whalen M, Stall R and Staskawicz J (1999). Expression of the Bs2 pepper gene confers resistance to bacteria spot disease in tomato. *Proc Natl Acad Sci USA*. 96, 14153-14158.
- The Arabidopsis genome initiative (2000). Analysis of the genome sequence of the flowering plant *Arabidopsis thaliana*. *Nature*. 408, 796-815.
- Thomas C, Jones D, Parniske M, Harrison K, Balint-Kurti P et al. (1997). Characterization of the tomato Cf-4 gene for resistance to *Cladosporium fulvum* identifies sequences that determine recognitional specificity in Cf-4 and Cf-9. *Plant Cell*. 9, 2209-2224.
- Tornero P, Merritt P, Sadanandom A, Shirasu K, Innes R, and Dangl J (2002). RAR1 and NDR1 contribute quantitatively to disease resistance in *Arabidopsis*, and their relative contributions are dependent on the R gene assayed. *The plant cell*. 14: 1005-1015
- Van Der Biezen E and Jones D (1998). Plant disease-resistance proteins and the gene-for-gene concept. *Trend in Biochemistry Science*. 23, 454-456.
- Van der Hoorn R, De Wit P and Joosten M (2002). Balancing selection favors guarding resistance proteins. *Trends in plant science*. 7: 67-71
- Van der Vossen E, Rouppe van der Voort J, Kanyuka K, Bendahmane A, Sandbrink H et al (2000). Homologue of a single resistance-gene cluster in potato confer resistance to distinct pathogens: a virus and a nematode. *Plant Journal*. 23: 567-576.
- Wang G, Ruan D, Song W, Sideris S, Chen L, Pi L, Zhang S, Zhang Z, Fauquet C, Gaut B, Ronald P (1998). Xa21D encodes a receptor-like molecule with a

- leucine-rich repeat domain that determines race specific recognition and is subject to adaptive evolution. *Plant Cell*. 10, 1-15
- Warren R, Henk A, Mowery P, Holub E, Innes R (1998).** A mutation within the leucine-rich repeat domain of the *Arabidopsis* disease resistance gene RPS5 partially suppresses multiple bacterial and downy mildew resistance genes. *Plant Cell*. 10, 1439-1452.
- Wees S, Swart E, Pelt J, Loon L and Pieterse C (2000).** Enhancement of induced disease resistance by simultaneous activation of salicylate- and jasmonate-dependent defense pathways in *Arabidopsis thaliana*. *Proc Natl Acad Sci USA*. 97, 8711-8716.
- Whitham S (1996).** The N gene of tobacco confers resistance to tobacco mosaic virus in transgenic tomato. *Proc Natl Acad Sci USA*. 93, 8776-8781.
- Yoshimura S, Yamanouchi U, Katayose Y, Toki S, Z-X et al (1998).** Expression of Xa1, a bacterial blight-resistance gene in rice, is induced by bacterial inoculation. *PNAS*. 95, 1663-1668.
- Yu D, Chen C and Chen Z (2001).** Evidence for an important role of WRKY DNA binding proteins in the regulation of NPR1 gene expression. *Plant Cell*. 13: 1527-1540.
- Yu Y, Buss G and Maroof M (1996).** Isolation of a superfamily of candidate disease-resistance genes in soybean based on a conserved nucleotide-binding site. *Proc Natl Acad Sci USA*. 93, 11751-11756.
- Zhang Y, Fan W, Kinkema M, Li X and Dong X (1999).** Interaction of NPR1 with basic leucine zipper protein transcription factors that bind sequences required for salicylic acid and induction of PR-1 gene. *Proc Natl Acad Sci USA*. 96, 6523-6528.
- Zou H, Henzel W, Liu X, Lutschg A and Wang X (1997).** Apaf-1, a human protein homologous to *C. elegans* CED-4, participates in Cytochrome c-dependent activation of caspase -3. *Cell*. 90: 405-413.
- Zhou J, Trifa Y, Silva, Pontier D, Lam E, Shah J, Klessig D (2000).** NPR1 differentially interacts with members of the TGA/OBF family of transcription

factors that bind an element of the PR-1 gene required for induction by salicylic acid. *Mol Plant Microbe Interact.* 13, 191-202.

SEQUENCE LISTING

<110> Queensland University of Technology
 <120> Resistance elements and uses therefor
 <130> 12177852
 <140> Unassigned
 <141> 2003-09-25
 <160> 4
 <170> PatentIn version 3.2
 <210> 1
 <211> 4380
 <212> DNA
 <213> Musa acuminata

<220>
 <221> CDS
 <222> (1)..(4323)

<400> 1
 atg tcg acg gcg cta gta atc gga gga tgg ttc gcg caa agc ttc atc 48
 Met Ser Thr Ala Leu Val Ile Gly Gly Trp Phe Ala Gln Ser Phe Ile
 1 5 10 15
 cag acg ttg ctc gac aag gcc agc aac tgc gcg atc caa caa ctc gcg 96
 Gln Thr Leu Leu Asp Lys Ala Ser Asn Cys Ala Ile Gln Gln Leu Ala
 20 25 30
 cgg cgc cgc ggc ctt cac gat gac ctg agg cgg ctg cgg acg tct ctg 144
 Arg Arg Arg Gly Leu His Asp Asp Leu Arg Arg Leu Arg Thr Ser Leu
 35 40 45
 ctc cgg atc cat gcc atc ctc gac aag gca gag acg agg tgg aac cat 192
 Leu Arg Ile His Ala Ile Leu Asp Lys Ala Glu Thr Arg Trp Asn His
 50 55 60
 aaa aac acg agc ttg gtg gag ctg gtg agg cag ctc aag gat gct gcc 240
 Lys Asn Thr Ser Leu Val Glu Leu Val Arg Gln Leu Lys Asp Ala Ala
 65 70 75 80
 tat gac gcc gag gac tta ctg gag gag ttg gag tac caa gcc gcg aag 288
 Tyr Asp Ala Glu Asp Leu Leu Glu Glu Leu Glu Tyr Gln Ala Ala Lys
 85 90 95
 caa aag gtc gag cac cgg gga gac cag ata agc gac ctc ttt tct ttt 336
 Gln Lys Val Glu His Arg Gly Asp Gln Ile Ser Asp Leu Phe Ser Phe
 100 105 110
 tcc ctt agt act gcg agc gag tgg ttg ggt gcc gat ggt gat gat gct 384
 Ser Leu Ser Thr Ala Ser Glu Trp Leu Gly Ala Asp Gly Asp Asp Ala
 115 120 125
 ggg act cga ttg agg gag atc cag ggg aag ctg tgc aac att gct gcc 432
 Gly Thr Arg Leu Arg Glu Ile Gln Gly Lys Leu Cys Asn Ile Ala Ala
 130 135 140

gat atg atg gat gtc atg cag cta ttg gca ccc gat gat ggg ggg aga Asp Met Met Asp Val Met Gln Leu Leu Ala Pro Asp Asp Gly Gly Arg 145 150 155 160	480
caa ttc gac tgg aag gtg gtg aga aga gaa acg agc tct ttc ttg acc Gln Phe Asp Trp Lys Val Val Arg Arg Glu Thr Ser Ser Phe Leu Thr 165 170 175	528
gaa acc gtc gtg ttt ggt cgg gac caa gaa agg gag aaa gta gta gaa Glu Thr Val Val Phe Gly Arg Asp Gln Glu Arg Glu Lys Val Val Glu 180 185 190	576
ttg ctg ttg gat tca gga tct ggt aac agt agc ttc tct gtc tta ccc Leu Leu Leu Asp Ser Gly Ser Gly Asn Ser Ser Phe Ser Val Leu Pro 195 200 205	624
ctc gtc gga atc gga ggg gtt ggg aag acg act ctg gct cag ctc gtg Leu Val Gly Ile Gly Gly Val Gly Lys Thr Thr Leu Ala Gln Leu Val 210 215 220	672
tac aac gac aat cgt gtc ggc aac tat ttc cac ctc aag gtt tgg gtc Tyr Asn Asp Asn Arg Val Gly Asn Tyr Phe His Leu Lys Val Trp Val 225 230 235 240	720
tgt gta tcc gac aat ttc aat gtg aag aga ctg acc aaa gag ata atc Cys Val Ser Asp Asn Phe Asn Val Lys Arg Leu Thr Lys Glu Ile Ile 245 250 255	768
gag tct gct acc aag gtg gaa caa tct gac aaa ttg aac ttg gac acc Glu Ser Ala Thr Lys Val Glu Gln Ser Asp Lys Leu Asn Leu Asp Thr 260 265 270	816
ctg caa cag atc ctc aag gag aag att gct tca gag agg ttt ctg cta Leu Gln Gln Ile Leu Lys Glu Lys Ile Ala Ser Glu Arg Phe Leu Leu 275 280 285	864
gtc ctc gat gat gtg tgg agc gaa aac agg gat gac tgg gaa agg ctg Val Leu Asp Asp Val Trp Ser Glu Asn Arg Asp Asp Trp Glu Arg Leu 290 295 300	912
tgc gca cca cta agg ttt gca gca aga ggc agc aag gtt ata gtc aca Cys Ala Pro Leu Arg Phe Ala Ala Arg Gly Ser Lys Val Ile Val Thr 305 310 315 320	960
act cga gac aca aag att gcc agc atc att ggc aca atg aag gaa att Thr Arg Asp Thr Lys Ile Ala Ser Ile Ile Gly Thr Met Lys Glu Ile 325 330 335	1008
tcg ctc gat ggt ctc cag gat gat gct tac tgg gag ctg ttc aag aaa Ser Leu Asp Gly Leu Gln Asp Asp Ala Tyr Trp Glu Leu Phe Lys Lys 340 345 350	1056
tgt gca ttt ggt tct gtg aac ccc cag gag cat cta gag ctc gag gtt Cys Ala Phe Gly Ser Val Asn Pro Gln Glu His Leu Glu Leu Glu Val 355 360 365	1104
atc ggt aga aag att gct ggt aag ttg aag ggc tca ccg cta gca gca Ile Gly Arg Lys Ile Ala Gly Lys Leu Lys Gly Ser Pro Leu Ala Ala 370 375 380	1152
aaa aca cta gga agc ttg ttg cgg ttg gat gtc agc caa gaa cac tgg	1200

- 3 -

ggc tgt caa tta cgg agt ttc cca caa ggc atg agc aag ctg atc aac Gly Cys Gln Leu Arg Ser Phe Pro Gln Gly Met Ser Lys Leu Ile Asn 645 650 655	1968
ttg agg caa ctt cgt gta gaa gat gag ata att tcc aag ata tat gag Leu Arg Gln Lys Leu Arg Val Glu Asp Glu Ile Ile Ser Lys Ile Tyr Glu 660 665 670	2016
gtt ggg aag ctg att tct ctg caa gaa ttg tct gca ttc aaa gtg cta Val Gly Lys Leu Ile Ser Leu Gln Glu Leu Ser Ala Phe Lys Val Leu 675 680 685	2064
aat aat cat gga aac aaa ctt gca gaa cta agt ggt ttg aca caa ctc Asn Asn His Gly Asn Lys Leu Ala Glu Leu Ser Gly Leu Thr Gln Leu 690 695 700	2112
cgc agc act cta cga att aca aat ctt gaa aat gta ggg agt aaa gaa Arg Ser Thr Leu Arg Ile Thr Asn Leu Glu Asn Val Gly Ser Lys Glu 705 710 715 720	2160
gaa gca agc aag gct aaa ctg cac agg aaa cag tat ctt gaa gca tta Glu Ala Ser Lys Ala Lys Leu His Arg Lys Gln Tyr Leu Glu Ala Leu 725 730 735	2208
gag tta gag tgg gca gct ggc cag gtt tcc agc ttg gag cat gag tta Glu Leu Glu Trp Ala Ala Gly Gln Val Ser Ser Leu Glu His Glu Leu 740 745 750	2256
ctt gtc tcg gag gaa gta ctt tta ggt ctc caa cca cat cac ttc ctc Leu Val Ser Glu Glu Val Leu Leu Gly Leu Gln Pro His His Phe Leu 755 760 765	2304
aaa agt ttg aca atc aga ggg tac agt ggt gca aca gta ccc agt tgg Lys Ser Leu Thr Ile Arg Gly Tyr Ser Gly Ala Thr Val Pro Ser Trp 770 775 780	2352
ctg gat gtg aaa atg cta ccg aac ttg gga act ctt aaa cta gag aac Leu Asp Val Lys Met Leu Pro Asn Leu Gly Thr Leu Lys Leu Glu Asn 785 790 795 800	2400
tgt aca aga ctg gag ggt ctt tca tat att gga caa ctg cca cat ctc Cys Thr Arg Leu Glu Gly Leu Ser Tyr Ile Gly Gln Leu Pro His Leu 805 810 815	2448
aag gtc ctt cat atg aag aga atg cct gtg gtg aaa caa atg agt cat Lys Val Leu His Met Lys Arg Met Pro Val Val Lys Gln Met Ser His 820 825 830	2496
gaa tta tgt ggc tgt acg aaa agc aag ttg ttc cct agg cta gaa gag Glu Leu Cys Gly Cys Thr Lys Ser Lys Leu Phe Pro Arg Leu Glu Glu 835 840 845	2544
tta gta ctg gag gat atg cca aca ttg aaa gaa ttc ccg aat ctt gca Leu Val Leu Glu Asp Met Pro Thr Leu Lys Glu Phe Pro Asn Leu Ala 850 855 860	2592
caa ctt cct tgt ctc aag att att cac atg aag aac atg ttt gca gta Gln Leu Pro Cys Leu Lys Ile Ile His Met Lys Asn Met Phe Ala Val 865 870 875 880	2640
aaa cat ata ggt cgt gaa tta tat ggt gat ata gag agc aat tgt ttt	2688

Lys His Ile Gly Arg Glu Leu Tyr Gly Asp Ile Glu Ser Asn Cys Phe	
885	890 895
cta tca tta gaa gag ctt gtg ctg cag gac atg ctg aca ttg gag gaa	2736
Leu Ser Leu Glu Glu Leu Val Leu Gln Asp Met Leu Thr Leu Glu Glu	
900	905 910
ctc cca aat ctt gga caa ctt cca cat ctt aag gtt att cac atg aag	2784
Leu Pro Asn Leu Gly Gln Leu Pro His Leu Lys Val Ile His Met Lys	
915	920 925
aac atg tct gca ctg aaa ctt ata ggt cgt gaa tta tgt gat tct aga	2832
Asn Met Ser Ala Leu Lys Leu Ile Gly Arg Glu Leu Cys Asp Ser Arg	
930	935 940
gag aaa att tgg ttt cct agg cta gaa gtg cta gtg ctg aag aac atg	2880
Glu Lys Ile Trp Phe Pro Arg Leu Glu Val Leu Val Leu Lys Asn Met	
945	950 955 960
ctg gca ctg gag gaa ctc cca agc ttg gac aac ttc cgt gtc tca aga	2928
Leu Ala Leu Glu Glu Leu Pro Ser Leu Asp Asn Phe Arg Val Ser Arg	
965	970 975
ttc ttc gca tcc agt gtc gaa gta ggc cat gga ctc ttt agt gct acg	2976
Phe Phe Ala Ser Val Glu Val Gly His Gly Leu Phe Ser Ala Thr	
980	985 990
agg aat aaa tgg ttt cca agg ctg gaa gag cta gaa atc aag ggc atg	3024
Arg Asn Lys Trp Phe Pro Arg Leu Glu Glu Leu Glu Ile Lys Gly Met	
995	1000 1005
ctg aca ttt gag gaa ctc cat tot ctt gaa aaa ctg cca tgt ctc	3069
Leu Thr Phe Glu Glu Leu His Ser Leu Glu Lys Leu Pro Cys Leu	
1010	1015 1020
aag gtt ttc cgc atc aag gga ttg cca gca gtg aaa aag ata ggc	3114
Lys Val Phe Arg Ile Lys Gly Leu Pro Ala Val Lys Lys Ile Gly	
1025	1030 1035
cat gga tta ttt gat tct acc tgt cag aga gag tgt ttt cca agg	3159
His Gly Leu Phe Asp Ser Thr Cys Gln Arg Glu Cys Phe Pro Arg	
1040	1045 1050
ttg gaa gat ctc gta tta agc gac atg cca gca tgg gaa gag tgg	3204
Leu Glu Asp Leu Val Leu Ser Asp Met Pro Ala Trp Glu Glu Trp	
1055	1060 1065
tcg tgg gct gaa agg gag gag tta ttt tcc tgc ttg tgt aga ctt	3249
Ser Trp Ala Glu Arg Glu Glu Leu Phe Ser Cys Leu Cys Arg Leu	
1070	1075 1080
aaa att gaa caa tgc ccc aaa ctt aaa tgc ttg ctt ccc atc cct	3294
Lys Ile Glu Gln Cys Pro Lys Leu Lys Cys Leu Leu Pro Ile Pro	
1085	1090 1095
cat tct ctc ata aaa ctt gaa tta tgg caa gtt ggg ctg aca gga	3339
His Ser Leu Ile Lys Leu Glu Leu Trp Gln Val Gly Leu Thr Gly	
1100	1105 1110
ctt cca gga tta tgc aaa gga att ggt gga ggt agc agc act aga	3384
Leu Pro Gly Leu Cys Lys Gly Ile Gly Gly Gly Ser Ser Thr Arg	
1115	1120 1125

act gct	tct ctc tca ctc ttg	cac att att aaa tgt	cca aat ctg	3429
Thr Ala	Ser Leu Ser Leu Leu	His Ile Ile Lys Cys	Pro Asn Leu	
1130	1135	1140		
aga aat	ctg gga gaa ggg ttg	cta tca aac cac ctg	cca cat atc	3474
Arg Asn	Leu Gly Glu Gly Leu	Leu Ser Asn His Leu	Pro His Ile	
1145	1150	1155		
aat gct	att cgg ata tgg gaa	tgt gct gaa ctg ttg	tgg ctg cct	3519
Asn Ala	Ile Arg Ile Trp Glu	Cys Ala Glu Leu Leu	Trp Leu Pro	
1160	1165	1170		
gtc aag	agg ttt aga gaa ttc	acc acc ctt gag aac	ttg tca ata	3564
Val Lys	Arg Phe Arg Glu Phe	Thr Thr Leu Glu Asn	Leu Ser Ile	
1175	1180	1185		
agg aac	tgc ccc aag ctc atg	agc atg aca cag tgt	gag gag aat	3609
Arg Asn	Cys Pro Lys Leu Met	Ser Met Thr Gln Cys	Glu Glu Asn	
1190	1195	1200		
gac ctc	ctc ctc ccg ccg tta	atc aag gca cta gaa	ttg ggt gac	3654
Asp Leu	Leu Leu Pro Pro Leu	Ile Lys Ala Leu Glu	Leu Gly Asp	
1205	1210	1215		
tgt gga	aat ctt ggg aaa tgc	ctg cct gga tgc cta	cac aac ctc	3699
Cys Gly	Asn Leu Gly Lys Ser	Leu Pro Gly Cys Leu	His Asn Leu	
1220	1225	1230		
agc tca	ctt act cag ttg gcg	ata tcc aat tgt cca	tac atg gta	3744
Ser Ser	Leu Thr Gln Leu Ala	Ile Ser Asn Cys Pro	Tyr Met Val	
1235	1240	1245		
tcc ctt	cca agg gaa gta atg	ctt cac ttg aag gaa	ctt gga act	3789
Ser Leu	Pro Arg Glu Val Met	Leu His Leu Lys Glu	Leu Gly Thr	
1250	1255	1260		
gta agg	atc gag aat tgt gat	ggg ctg gga tca ata	gag ggt tta	3834
Val Arg	Ile Glu Asn Cys Asp	Gly Leu Gly Ser Ile	Glu Gly Leu	
1265	1270	1275		
caa gtt	ctc aaa tca ctc aag	aga ttg gca atc ata	gga tgt ccc	3879
Gln Val	Leu Lys Ser Leu Lys	Arg Leu Ala Ile Ile	Gly Cys Pro	
1280	1285	1290		
agg ctt	ttg cta aat gaa ggg	gat gag caa ggg gag	gtc ttg tca	3924
Arg Leu	Leu Leu Asn Glu Gly	Asp Glu Gln Gly Glu	Val Leu Ser	
1295	1300	1305		
ctg ctt	gaa tta tca gta gat	aaa aca gcc cta ctt	aaa ctc tca	3969
Leu Leu	Glu Leu Ser Val Asp	Lys Thr Ala Leu Leu	Lys Leu Ser	
1310	1315	1320		
ctt ata	aaa aat aca cta cca	ttc atc cat tct ctc	aga atc atc	4014
Leu Ile	Lys Asn Thr Leu Pro	Phe Ile His Ser Leu	Arg Ile Ile	
1325	1330	1335		
tgg tct	cct cag aaa gtg atg	ttt gac ttg gag gag	cag gaa ttg	4059
Trp Ser	Pro Gln Lys Val Met	Phe Asp Leu Glu Glu	Gln Glu Leu	
1340	1345	1350		
gtg cac	agc ctc aca gct ctc	agg cgc ctt gaa ttc	ttc aga tgc	4104

Val His Ser Leu Thr Ala Leu Arg Arg Leu Glu Phe Phe Arg Cys
 1355 1360 1365
 aag aat ctc cag tcc ttg cca aca gag ttg cat acc ctt cct tcc 4149
 Lys Asn Leu Gln Ser Leu Pro Thr Glu Leu His Thr Leu Pro Ser
 1370 1375 1380
 ctc cat gct ttg gtt gta agt gac tgc cca cag atc caa tca ctg 4194
 Leu His Ala Leu Val Val Ser Asp Cys Pro Gln Ile Gln Ser Leu
 1385 1390 1395
 ccg gag aag gga ctc ccg aca ctc ctc aca gat tta gga ttt gac 4239
 Pro Glu Lys Gly Leu Pro Thr Leu Leu Thr Asp Leu Gly Phe Asp
 1400 1405 1410
 cat tgc cac cca gtg ctg act gcg caa ctg gaa aag cac ctg gca 4284
 His Cys His Pro Val Leu Thr Ala Gln Leu Glu Lys His Leu Ala
 1415 1420 1425
 gag atg aag agc tca ggt cga ttt cac cca gtt tat gca taggcaacat 4333
 Glu Met Lys Ser Ser Gly Arg Phe His Pro Val Tyr Ala
 1430 1435 1440
 gagtgaggat ggagaaaggg gagtggaaga gaaagatttc gattgcc 4380

 <210> 2
 <211> 1441
 <212> PRT
 <213> Musa acuminata

 <400> 2
 Met Ser Thr Ala Leu Val Ile Gly Gly Trp Phe Ala Gln Ser Phe Ile
 1 5 10 15
 Gln Thr Leu Leu Asp Lys Ala Ser Asn Cys Ala Ile Gln Gln Leu Ala
 20 25 30
 Arg Arg Arg Gly Leu His Asp Asp Leu Arg Arg Leu Arg Thr Ser Leu
 35 40 45
 Leu Arg Ile His Ala Ile Leu Asp Lys Ala Glu Thr Arg Trp Asn His
 50 55 60
 Lys Asn Thr Ser Leu Val Glu Leu Val Arg Gln Leu Lys Asp Ala Ala
 65 70 75 80
 Tyr Asp Ala Glu Asp Leu Leu Glu Glu Leu Glu Tyr Gln Ala Ala Lys
 85 90 95
 Gln Lys Val Glu His Arg Gly Asp Gln Ile Ser Asp Leu Phe Ser Phe
 100 105 110
 Ser Leu Ser Thr Ala Ser Glu Trp Leu Gly Ala Asp Gly Asp Asp Ala

115

120

125

Gly Thr Arg Leu Arg Glu Ile Gln Gly Lys Leu Cys Asn Ile Ala Ala
130 135 140

Asp Met Met Asp Val Met Gln Leu Leu Ala Pro Asp Asp Gly Gly Arg
145 150 155 160

Gln Phe Asp Trp Lys Val Val Arg Arg Glu Thr Ser Ser Phe Leu Thr
165 170 175

Glu Thr Val Val Phe Gly Arg Asp Gln Glu Arg Glu Lys Val Val Glu
180 185 190

Leu Leu Leu Asp Ser Gly Ser Gly Asn Ser Ser Phe Ser Val Leu Pro
195 200 205

Leu Val Gly Ile Gly Gly Val Gly Lys Thr Thr Leu Ala Gln Leu Val
210 215 220

Tyr Asn Asp Asn Arg Val Gly Asn Tyr Phe His Leu Lys Val Trp Val
225 230 235 240

Cys Val Ser Asp Asn Phe Asn Val Lys Arg Leu Thr Lys Glu Ile Ile
245 250 255

Glu Ser Ala Thr Lys Val Glu Gln Ser Asp Lys Leu Asn Leu Asp Thr
260 265 270

Leu Gln Gln Ile Leu Lys Glu Lys Ile Ala Ser Glu Arg Phe Leu Leu
275 280 285

Val Leu Asp Asp Val Trp Ser Glu Asn Arg Asp Asp Trp Glu Arg Leu
290 295 300

Cys Ala Pro Leu Arg Phe Ala Ala Arg Gly Ser Lys Val Ile Val Thr
305 310 315 320

Thr Arg Asp Thr Lys Ile Ala Ser Ile Ile Gly Thr Met Lys Glu Ile
325 330 335

Ser Leu Asp Gly Leu Gln Asp Asp Ala Tyr Trp Glu Leu Phe Lys Lys
340 345 350

Cys Ala Phe Gly Ser Val Asn Pro Gln Glu His Leu Glu Leu Glu Val
355 360 365

Ile Gly Arg Lys Ile Ala Gly Lys Leu Lys Gly Ser Pro Leu Ala Ala
370 375 380

Lys Thr Leu Gly Ser Leu Leu Arg Leu Asp Val Ser Gln Glu His Trp
385 390 395 400

Arg Thr Ile Met Glu Ser Glu Val Trp Gln Leu Pro Gln Ala Glu Asn
405 410 415

Glu Ile Leu Pro Val Leu Trp Leu Ser Tyr Gln His Leu Pro Gly His
420 425 430

Leu Arg Gln Cys Phe Ala Phe Cys Ala Val Phe His Lys Asp Tyr Leu
435 440 445

Phe Tyr Lys His Glu Leu Ile Gln Thr Trp Ile Ala Glu Gly Phe Ile
450 455 460

Ala His Gln Gly Asn Lys Arg Met Glu Asp Val Gly Ser Ser Tyr Phe
465 470 475 480

His Glu Leu Val Asn Arg Ser Phe Phe Gln Glu Ser Arg Trp Arg Gly
485 490 495

Arg Tyr Val Met His Asp Leu Ile His Asp Leu Ala Gln Phe Ile Ser
500 505 510

Val Gly Glu Cys His Arg Ile Asp Asp Asp Lys Ser Lys Glu Thr Pro
515 520 525

Ser Thr Thr Arg His Leu Ser Val Ala Leu Thr Glu Gln Met Lys Leu
530 535 540

Val Asp Phe Ser Gly Tyr Asn Lys Leu Arg Thr Leu Met Ile Asn Asn
545 550 555 560

Gln Arg Asn Gln Tyr Pro Tyr Met Thr Lys Val Asn Ser Cys Leu Leu
565 570 575

Pro His Ser Leu Phe Lys Arg Leu Lys Arg Ile His Val Leu Val Leu
580 585 590

Gln Lys Cys Gly Met Lys Glu Leu Pro Asp Ile Ile Gly Asp Leu Ile
595 600 605

Gln Leu Arg Tyr Leu Asp Ile Ser Tyr Asn Ala Cys Ile Gln Arg Leu

25 September 2003

610

615

620

Pro Glu Ser Leu Cys Asp Leu Tyr Asn Leu Gln Ala Leu Arg Leu Trp
625 630 635 640

Gly Cys Gln Leu Arg Ser Phe Pro Gln Gly Met Ser Lys Leu Ile Asn
645 650 655

Leu Arg Gln Leu Arg Val Glu Asp Glu Ile Ile Ser Lys Ile Tyr Glu
660 665 670

Val Gly Lys Leu Ile Ser Leu Gln Glu Leu Ser Ala Phe Lys Val Leu
675 680 685

Asn Asn His Gly Asn Lys Leu Ala Glu Leu Ser Gly Leu Thr Gln Leu
690 695 700

Arg Ser Thr Leu Arg Ile Thr Asn Leu Glu Asn Val Gly Ser Lys Glu
705 710 715 720

Glu Ala Ser Lys Ala Lys Leu His Arg Lys Gln Tyr Leu Glu Ala Leu
725 730 735

Glu Leu Glu Trp Ala Ala Gly Gln Val Ser Ser Leu Glu His Glu Leu
740 745 750

Leu Val Ser Glu Glu Val Leu Leu Gly Leu Gln Pro His His Phe Leu
755 760 765

Lys Ser Leu Thr Ile Arg Gly Tyr Ser Gly Ala Thr Val Pro Ser Trp
770 775 780

Leu Asp Val Lys Met Leu Pro Asn Leu Gly Thr Leu Lys Leu Glu Asn
785 790 795 800

Cys Thr Arg Leu Glu Gly Leu Ser Tyr Ile Gly Gln Leu Pro His Leu
805 810 815

Lys Val Leu His Met Lys Arg Met Pro Val Val Lys Gln Met Ser His
820 825 830

Glu Leu Cys Gly Cys Thr Lys Ser Lys Leu Phe Pro Arg Leu Glu Glu
835 840 845

Leu Val Leu Glu Asp Met Pro Thr Leu Lys Glu Phe Pro Asn Leu Ala
850 855 860

Gln Leu Pro Cys Leu Lys Ile Ile His Met Lys Asn Met Phe Ala Val
 865 870 875 880

Lys His Ile Gly Arg Glu Leu Tyr Gly Asp Ile Glu Ser Asn Cys Phe
 885 890 895

Leu Ser Leu Glu Glu Leu Val Leu Gln Asp Met Leu Thr Leu Glu Glu
 900 905 910

Leu Pro Asn Leu Gly Gln Leu Pro His Leu Lys Val Ile His Met Lys
 915 920 925

Asn Met Ser Ala Leu Lys Leu Ile Gly Arg Glu Leu Cys Asp Ser Arg
 930 935 940

Glu Lys Ile Trp Phe Pro Arg Leu Glu Val Leu Val Leu Lys Asn Met
 945 950 955 960

Leu Ala Leu Glu Glu Leu Pro Ser Leu Asp Asn Phe Arg Val Ser Arg
 965 970 975

Phe Phe Ala Ser Ser Val Glu Val Gly His Gly Leu Phe Ser Ala Thr
 980 985 990

Arg Asn Lys Trp Phe Pro Arg Leu Glu Glu Leu Glu Ile Lys Gly Met
 995 1000 1005

Leu Thr Phe Glu Glu Leu His Ser Leu Glu Lys Leu Pro Cys Leu
 1010 1015 1020

Lys Val Phe Arg Ile Lys Gly Leu Pro Ala Val Lys Lys Ile Gly
 1025 1030 1035

His Gly Leu Phe Asp Ser Thr Cys Gln Arg Glu Cys Phe Pro Arg
 1040 1045 1050

Leu Glu Asp Leu Val Leu Ser Asp Met Pro Ala Trp Glu Glu Trp
 1055 1060 1065

Ser Trp Ala Glu Arg Glu Glu Leu Phe Ser Cys Leu Cys Arg Leu
 1070 1075 1080

Lys Ile Glu Gln Cys Pro Lys Leu Lys Cys Leu Leu Pro Ile Pro
 1085 1090 1095

His Ser Leu Ile Lys Leu Glu Leu Trp Gln Val Gly Leu Thr Gly

1100	1105	1110
Leu Pro Gly Leu Cys Lys Gly Ile Gly Gly Gly Ser Ser Thr Arg 1115 1120 1125		
Thr Ala Ser Leu Ser Leu Leu His Ile Ile Lys Cys Pro Asn Leu 1130 1135 1140		
Arg Asn Leu Gly Glu Gly Leu Leu Ser Asn His Leu Pro His Ile 1145 1150 1155		
Asn Ala Ile Arg Ile Trp Glu Cys Ala Glu Leu Leu Trp Leu Pro 1160 1165 1170		
Val Lys Arg Phe Arg Glu Phe Thr Thr Leu Glu Asn Leu Ser Ile 1175 1180 1185		
Arg Asn Cys Pro Lys Leu Met Ser Met Thr Gln Cys Glu Glu Asn 1190 1195 1200		
Asp Leu Leu Leu Pro Pro Leu Ile Lys Ala Leu Glu Leu Gly Asp 1205 1210 1215		
Cys Gly Asn Leu Gly Lys Ser Leu Pro Gly Cys Leu His Asn Leu 1220 1225 1230		
Ser Ser Leu Thr Gln Leu Ala Ile Ser Asn Cys Pro Tyr Met Val 1235 1240 1245		
Ser Leu Pro Arg Glu Val Met Leu His Leu Lys Glu Leu Gly Thr 1250 1255 1260		
Val Arg Ile Glu Asn Cys Asp Gly Leu Gly Ser Ile Glu Gly Leu 1265 1270 1275		
Gln Val Leu Lys Ser Leu Lys Arg Leu Ala Ile Ile Gly Cys Pro 1280 1285 1290		
Arg Leu Leu Leu Asn Glu Gly Asp Glu Gln Gly Glu Val Leu Ser 1295 1300 1305		
Leu Leu Glu Leu Ser Val Asp Lys Thr Ala Leu Leu Lys Leu Ser 1310 1315 1320		
Leu Ile Lys Asn Thr Leu Pro Phe Ile His Ser Leu Arg Ile Ile 1325 1330 1335		

25 September 2003

Trp Ser Pro Gln Lys Val Met Phe Asp Leu Glu Glu Gln Glu Leu
 1340 1345 1350

Val His Ser Leu Thr Ala Leu Arg Arg Leu Glu Phe Phe Arg Cys
 1355 1360 1365

Lys Asn Leu Gln Ser Leu Pro Thr Glu Leu His Thr Leu Pro Ser
 1370 1375 1380

Leu His Ala Leu Val Val Ser Asp Cys Pro Gln Ile Gln Ser Leu
 1385 1390 1395

Pro Glu Lys Gly Leu Pro Thr Leu Leu Thr Asp Leu Gly Phe Asp
 1400 1405 1410

His Cys His Pro Val Leu Thr Ala Gln Leu Glu Lys His Leu Ala
 1415 1420 1425

Glu Met Lys Ser Ser Gly Arg Phe His Pro Val Tyr Ala
 1430 1435 1440

<210> 3
 <211> 3699
 <212> DNA
 <213> Musa acuminata spp malaccensis

<220>
 <221> CDS
 <222> (1)..(3696)

<400> 3
 atg gct gat gtc aca cca cag gca gcg gcg gtg ttc tcc ctg gtg aat 48
 Met Ala Asp Val Thr Pro Gln Ala Ala Val Phe Ser Leu Val Asn
 1 5 10 15
 gaa atc ttt aac cgg tcc atc aat ttg atc gtc gcg gaa ctc cgg ttg 96
 Glu Ile Phe Asn Arg Ser Ile Asn Leu Ile Val Ala Glu Leu Arg Leu
 20 25 30
 cag ttg aat gcg aga gcc gag ctg aac aat ctg cag aga aca cta ttg 144
 Gln Leu Asn Ala Arg Ala Glu Leu Asn Asn Leu Gln Arg Thr Leu Leu
 35 40 45
 agg act cac tct ctg ctc gag gag gca aag gcg agg cgg atg act gac 192
 Arg Thr His Ser Leu Leu Glu Glu Ala Lys Ala Arg Arg Met Thr Asp
 50 55 60
 aag tct ctc gtg ctg tgg ctg atg gag ctc aag gaa tgg gcc tac gac 240
 Lys Ser Leu Val Leu Trp Leu Met Glu Leu Lys Glu Trp Ala Tyr Asp
 65 70 75 80
 gcc gac gac atc ctc gac gag tac gag gcc gca gca atc cga ctg aag 288
 Ala Asp Asp Ile Leu Asp Glu Tyr Glu Ala Ala Ile Arg Leu Lys

85	90	95	
gta aca cgc tgc acc ttc aaa cgt ctt atc gat cat gtg att ata aat Val Thr Arg Ser Thr Phe Lys Arg Leu Ile Asp His Val Ile Ile Asn 100 105 110			336
gtt cca tta gcg cac aaa gta gca gac atc agg aaa agg ttg aac ggg Val Pro Leu Ala His Lys Val Ala Asp Ile Arg Lys Arg Leu Asn Gly 115 120 125			384
gtc act ctt gag agg gag cta aat ctg ggt gcg ctg gaa ggg tgc cag Val Thr Leu Glu Arg Glu Leu Asn Leu Gly Ala Leu Glu Gly Ser Gln 130 135 140			432
ccg ctt gat tcc acg aaa aga ggt gtg acc act tct ctt ctg act gaa Pro Leu Asp Ser Thr Lys Arg Gly Val Thr Thr Ser Leu Leu Thr Glu 145 150 155 160			480
tct tgt att gtc ggg cga gct caa gat aag gag aat ttg att cgg ttg Ser Cys Ile Val Gly Arg Ala Gln Asp Lys Glu Asn Leu Ile Arg Leu 165 170 175			528
ctg ttg gag ccc agc gat ggg gcg gtt cct gtt gtt cct ata gtt gga Leu Leu Glu Pro Ser Asp Gly Ala Val Pro Val Val Pro Ile Val Gly 180 185 190			576
tta gga ggg gca ggg aag acg act ctg tct cag ctt atc ttt aat gac Leu Gly Gly Ala Gly Lys Thr Thr Leu Ser Gln Leu Ile Phe Asn Asp 195 200 205			624
aag aga gtg gag gag cat ttc cca ttg aga atg tgg gtg tgt gtg tct Lys Arg Val Glu Glu His Phe Pro Leu Arg Met Trp Val Cys Val Ser 210 215 220			672
gac gat ttt gat gtg aag aga att act aga gag atc aca gag tac gcc Asp Asp Phe Asp Val Lys Arg Ile Thr Arg Glu Ile Thr Glu Tyr Ala 225 230 235 240			720
acc aac gga agg ttc atg gat ctc acc aac ttg aat atg ctt caa gtt Thr Asn Gly Arg Phe Met Asp Leu Thr Asn Leu Asn Met Leu Gln Val 245 250 255			768
aat ctg aaa gag gag ata agg ggg acg aca ttt ttg ctt gtg ctg gat Asn Leu Lys Glu Glu Ile Arg Gly Thr Thr Phe Leu Leu Val Leu Asp 260 265 270			816
gat gtg tgg aac gaa gac ccc gtg aag tgg gaa agc ctg tta gcc cca Asp Val Trp Asn Glu Asp Pro Val Lys Trp Glu Ser Leu Leu Ala Pro 275 280 285			864
tta gat gcc gga gga cgg gga agc gtg gtc att gtg acg aca cag agc Leu Asp Ala Gly Gly Arg Gly Ser Val Val Ile Val Thr Gln Ser 290 295 300			912
aaa aag gtc gcc gat gtc acc ggc acg atg gag cca tac gtt ctc gag Lys Lys Val Ala Asp Val Thr Gly Thr Met Glu Pro Tyr Val Leu Glu 305 310 315 320			960
gag tta acg gag gat gac agt tgg tca ctc atc gag agt cac tcc ttc Glu Leu Thr Glu Asp Ser Trp Ser Leu Ile Glu Ser His Ser Phe 325 330 335			1008

```

agg gag gcg agc tgc tct agt aca aat cct aga atg gaa gag atc ggg      1056
Arg Glu Ala Ser Cys Ser Ser Thr Asn Pro Arg Met Glu Glu Ile Gly
          340                      345                      350

agg aag ata gcc aag aag atc agt ggc cta cct tac gga gca aca gca      1104
Arg Lys Ile Ala Lys Lys Ile Ser Gly Leu Pro Tyr Gly Ala Thr Ala
          355                      360                      365

atg ggg aga tat cta aga tct aag cac gga gaa agc agc tgg aga gaa      1152
Met Gly Arg Tyr Leu Arg Ser Lys His Gly Glu Ser Ser Trp Arg Glu
          370                      375                      380

gtc ttg gaa act gag act tgg gag atg cca ccg gct gca agt gat gtg      1200
Val Leu Glu Thr Glu Thr Trp Glu Met Pro Pro Ala Ala Ser Asp Val
          385                      390                      395                      400

tta tcc gct cta agg aga agt tac gac aat cta ccc cct cag ctg aag      1248
Leu Ser Ala Leu Arg Arg Ser Tyr Asp Asn Leu Pro Pro Gln Leu Lys
          405                      410                      415

ctc tgt ttt gcc ttc tgt gct ctg ttt aca aag ggc tac agg ttt cga      1296
Leu Cys Phe Ala Phe Cys Ala Leu Phe Thr Lys Gly Tyr Arg Phe Arg
          420                      425                      430

aag gat aca ctg atc cac atg tgg ata gct caa aat ttg att caa tca      1344
Lys Asp Thr Leu Ile His Met Trp Ile Ala Gln Asn Leu Ile Gln Ser
          435                      440                      445

aca gag tcg aaa aga tcg gag gac atg gca gaa gaa tgc ttt gat gat      1392
Thr Glu Ser Lys Arg Ser Glu Asp Met Ala Glu Glu Cys Phe Asp Asp
          450                      455                      460

ttg gtg tgc aga ttc ttc ttt cgg tac tcc tgg ggc aac tat gtg atg      1440
Leu Val Cys Arg Phe Phe Phe Arg Tyr Ser Trp Gly Asn Tyr Val Met
          465                      470                      475                      480

aat gac tca gtc cat gac ctc gct cga tgg gtt tca ttg gat gaa tat      1488
Asn Asp Ser Val His Asp Leu Ala Arg Trp Val Ser Leu Asp Glu Tyr
          485                      490                      495

ttt cga gca gat gaa gac tca cca ttg cat att tca aag cca att cgt      1536
Phe Arg Ala Asp Glu Asp Ser Pro Leu His Ile Ser Lys Pro Ile Arg
          500                      505                      510

cat ttg tca tgg tgc agt gaa aga ata acc aat gtt ctt gag gat aat      1584
His Leu Ser Trp Cys Ser Glu Arg Ile Thr Asn Val Leu Glu Asp Asn
          515                      520                      525

aac act ggt gga gat gct gtc aat ccg ctc agc agt ttg cgc act ctc      1632
Asn Thr Gly Gly Asp Ala Val Asn Pro Leu Ser Ser Leu Arg Thr Leu
          530                      535                      540

ctt ttc tta ggc caa tct gag ttc cgg tcg tat cat ctt ctt gat aga      1680
Leu Phe Leu Gly Gln Ser Glu Phe Arg Ser Tyr His Leu Leu Asp Arg
          545                      550                      555                      560

atg ttc agg atg ttg agc cga atc cgt gtt ttg gat ttc agc aac tgc      1728
Met Phe Arg Met Leu Ser Arg Ile Arg Val Leu Asp Phe Ser Asn Cys
          565                      570                      575

gtc ata aga aat ttg cct tct tcg gtt gga aat ctg aaa cat ctg cgt      1776
Val Ile Arg Asn Leu Pro Ser Ser Val Gly Asn Leu Lys His Leu Arg

```

580					585					590						
tac	ctg	ggc	ctg	tct	aat	acg	aga	att	caa	agg	ttg	ccg	gag	tct	gta	1824
Tyr	Leu	Gly	Leu	Ser	Asn	Thr	Arg	Ile	Gln	Arg	Leu	Pro	Glu	Ser	Val	
595					600					605						
aca	cgt	ctt	tgc	ctc	ctt	cag	aca	ttg	cta	cta	gag	ggc	tgt	gaa	ctg	1872
Thr	Arg	Leu	Cys	Leu	Leu	Gln	Thr	Leu	Leu	Leu	Glu	Gly	Cys	Glu	Leu	
610					615					620						
tgc	agg	tta	cca	aga	agc	atg	agc	agg	ctc	gtc	aaa	ctg	agg	cag	ctc	1920
Cys	Arg	Leu	Pro	Arg	Ser	Met	Ser	Arg	Leu	Val	Lys	Leu	Arg	Gln	Leu	
625					630					635					640	
aaa	gca	aat	cca	gat	gta	att	gcc	gac	ata	gcc	aaa	gtc	ggg	aga	ttg	1968
Lys	Ala	Asn	Pro	Asp	Val	Ile	Ala	Asp	Ile	Ala	Lys	Val	Gly	Arg	Leu	
645					650					655						
atc	gaa	ctt	caa	gag	ctg	aaa	gcc	tat	aat	gtt	gac	aag	aaa	aaa	gga	2016
Ile	Glu	Leu	Gln	Glu	Leu	Lys	Ala	Tyr	Asn	Val	Asp	Lys	Lys	Lys	Gly	
660					665					670						
cat	ggg	att	gca	gag	cta	agt	gca	atg	aat	cag	ctt	cac	ggg	gat	ctt	2064
His	Gly	Ile	Ala	Glu	Leu	Ser	Ala	Met	Asn	Gln	Leu	His	Gly	Asp	Leu	
675					680					685						
tcc	att	aga	aac	ctt	caa	aat	gta	gag	aaa	acg	cga	gag	tct	cgg	aag	2112
Ser	Ile	Arg	Asn	Leu	Gln	Asn	Val	Glu	Lys	Thr	Arg	Glu	Ser	Arg	Lys	
690					695					700						
gcg	agg	ttg	gac	gag	aaa	cag	aag	ctt	aag	ctc	ttg	gat	ctg	cga	tgg	2160
Ala	Arg	Leu	Asp	Glu	Lys	Gln	Lys	Leu	Lys	Leu	Leu	Asp	Leu	Arg	Trp	
705					710					715					720	
gct	gac	ggg	agg	ggg	gcc	gga	gaa	tgt	gat	cgt	gac	agg	aaa	gtt	ctt	2208
Ala	Asp	Gly	Arg	Gly	Ala	Gly	Glu	Cys	Asp	Arg	Asp	Arg	Lys	Val	Leu	
725					730					735						
aaa	ggc	ctc	cga	cca	cat	cca	aac	ctg	aga	gaa	ttg	agt	atc	aaa	tac	2256
Lys	Gly	Leu	Arg	Pro	His	Pro	Asn	Leu	Arg	Glu	Leu	Ser	Ile	Lys	Tyr	
740					745					750						
tac	gga	ggc	act	tca	tct	ccg	agt	tgg	atg	acg	gat	cag	tat	ctg	ccc	2304
Tyr	Gly	Gly	Thr	Ser	Ser	Pro	Ser	Trp	Met	Thr	Asp	Gln	Tyr	Leu	Pro	
755					760					765						
aac	atg	gaa	acg	att	cgc	ctg	cgt	agc	tgc	gca	agg	ttg	acg	gaa	ctc	2352
Asn	Met	Glu	Thr	Ile	Arg	Leu	Arg	Ser	Cys	Ala	Arg	Leu	Thr	Glu	Leu	
770					775					780						
cca	tgt	ctc	ggg	cag	ctg	cat	atc	ctt	aga	cat	ttg	cac	atc	gat	ggg	2400
Pro	Cys	Leu	Gly	Gln	Leu	His	Ile	Leu	Arg	His	Leu	His	Ile	Asp	Gly	
785					790					795					800	
atg	tcc	caa	gtg	aga	caa	att	aat	ctg	caa	ttt	tat	ggc	acc	gga	gaa	2448
Met	Ser	Gln	Val	Arg	Gln	Ile	Asn	Leu	Gln	Phe	Tyr	Gly	Thr	Gly	Glu	
805					810					815						
gtt	tca	ggg	ttt	cca	ttg	ctg	gag	ctc	ctg	aac	ata	cgt	cgc	atg	ccc	2496
Val	Ser	Gly	Phe	Pro	Leu	Leu	Glu	Leu	Leu	Asn	Ile	Arg	Arg	Met	Pro	
820					825					830						

agt ctg gag gaa tgg tgc gaa cca cgg aga aac tgt tgc tac ttc cct	2544
Ser Leu Glu Glu Trp Ser Glu Pro Arg Arg Asn Cys Cys Tyr Phe Pro	
835 840 845	
cgc ctc cat aaa ctg ctg atc gag gat tgt ccc agg ctc agg aat ctg	2592
Arg Leu His Lys Leu Leu Ile Glu Asp Cys Pro Arg Leu Arg Asn Leu	
850 855 860	
ccc tcc ctc cca cca aca ctg gaa gaa cta agg ata tca aga aca gga	2640
Pro Ser Leu Pro Pro Thr Leu Glu Glu Leu Arg Ile Ser Arg Thr Gly	
865 870 875 880	
cta gtt gat ctt cca gga ttc cat gga aac ggt gat gtg acg acg aat	2688
Leu Val Asp Leu Ser Gly Phe His Gly Asn Gly Asp Val Thr Thr Asn	
885 890 895	
gtt tcc ctt tct tct ttg cat gtt tgc gag tgt cga gaa ctg aga tcc	2736
Val Ser Leu Ser Ser Leu His Val Ser Glu Cys Arg Glu Leu Arg Ser	
900 905 910	
cta agc gaa gga ttg ttg cag cac aac ctc gtc gcc ctc aag aca gcg	2784
Leu Ser Glu Gly Leu Leu Gln His Asn Leu Val Ala Leu Lys Thr Ala	
915 920 925	
gca ttt acc gat tgt gat tct ctt gag ttt ttg ccg gcg gaa gga ttc	2832
Ala Phe Thr Asp Cys Asp Ser Leu Glu Phe Leu Pro Ala Glu Gly Phe	
930 935 940	
aga aca gcc att tca ctt gaa tca ttg ata atg act aat tgt cca ctg	2880
Arg Thr Ala Ile Ser Leu Glu Ser Leu Ile Met Thr Asn Cys Pro Leu	
945 950 955 960	
cct tgc agt ttt ctt ttg cct tcc tct ctc gag cat cta aag ttg cag	2928
Pro Cys Ser Phe Leu Leu Pro Ser Ser Leu Glu His Leu Lys Leu Gln	
965 970 975	
cca tgc ctc tat cca aac aac aat gag gat tca ctg tca aca tgc ttc	2976
Pro Cys Leu Tyr Pro Asn Asn Asn Glu Asp Ser Leu Ser Thr Cys Phe	
980 985 990	
gag aac ctc aca tct ctt tcc ttc ttg gac atc aaa gat tgt cca aat	3024
Glu Asn Leu Thr Ser Leu Ser Phe Leu Asp Ile Lys Asp Cys Pro Asn	
995 1000 1005	
ctg tca tca ttt cca ccg ggt cct cta tgt cag cta tca gca ctc	3069
Leu Ser Ser Phe Pro Pro Gly Pro Leu Cys Gln Leu Ser Ala Leu	
1010 1015 1020	
caa cat ttg tcc ctc gtc aat tgc cag agg cta caa tct att ggc	3114
Gln His Leu Ser Leu Val Asn Cys Gln Arg Leu Gln Ser Ile Gly	
1025 1030 1035	
ttc cag gca ctc acc tcc ctc gaa agc ttg aca att cag aac tgc	3159
Phe Gln Ala Leu Thr Ser Leu Glu Ser Leu Thr Ile Gln Asn Cys	
1040 1045 1050	
cct cgc ctc acc atg tca cac agt ttg gtt gag gtg aat aac tct	3204
Pro Arg Leu Thr Met Ser His Ser Leu Val Glu Val Asn Asn Ser	
1055 1060 1065	
tcc gat aca ggg ctc gcg ttt aat atc act cga tgg atg cgc aga	3249
Ser Asp Thr Gly Leu Ala Phe Asn Ile Thr Arg Trp Met Arg Arg	

1070	1075	1080	
cga aca ggt gac gac ggc ttg	atg ctc aga cac	cga gca caa aat	3294
Arg Thr Gly Asp Asp Gly Leu	Met Leu Arg His	Arg Ala Gln Asn	
1085	1090	1095	
gat tca ttt ttc ggg gga ctt	ctg caa cac ctc acc	ttc ctc cag	3339
Asp Ser Phe Phe Gly Gly Leu	Leu Gln His Leu Thr	Phe Leu Gln	
1100	1105	1110	
ttt cta aag atc tgc cag tgt	cca caa ctc gta acc	ttc acc ggc	3384
Phe Leu Lys Ile Cys Gln Cys	Pro Gln Leu Val Thr	Phe Thr Gly	
1115	1120	1125	
gaa gag gaa gag aag tgg aga	aac ctt act tct ctt	caa att ctg	3429
Glu Glu Glu Glu Lys Trp Arg	Asn Leu Thr Ser Leu	Gln Ile Leu	
1130	1135	1140	
cac atc gtt gat tgt cca aac	ctg gag gta ctg cct	gca aac ttg	3474
His Ile Val Asp Cys Pro Asn	Leu Glu Val Leu Pro	Ala Asn Leu	
1145	1150	1155	
caa agc ctc tgc tcc ctc agc	acc ttg tac atc gtc	aga tgc cca	3519
Gln Ser Leu Cys Ser Leu Ser	Thr Leu Tyr Ile Val	Arg Cys Pro	
1160	1165	1170	
aga atc cat gcg ttt cct ccc	gga ggt gtc agc atg	tcc ctg gca	3564
Arg Ile His Ala Phe Pro Pro	Gly Gly Val Ser Met	Ser Leu Ala	
1175	1180	1185	
cat ttg gtc atc cat gaa tgc	cct cag ctg tgt cag	cga tgt gat	3609
His Leu Val Ile His Glu Cys	Pro Gln Leu Cys Gln	Arg Cys Asp	
1190	1195	1200	
cca ccg gga ggt gat gat tgg	ccc tta ata gct aat	gta cca aga	3654
Pro Pro Gly Gly Asp Asp Trp	Pro Leu Ile Ala Asn	Val Pro Arg	
1205	1210	1215	
ata tgt ctt gga agg act cat	cca tgt cgc tgt agc	acc acc tga	3699
Ile Cys Leu Gly Arg Thr His	Pro Cys Arg Cys Ser	Thr Thr	
1220	1225	1230	
<210> 4			
<211> 1232			
<212> PRT			
<213> Musa acuminata spp malaccensis			
<400> 4			
Met Ala Asp Val Thr Pro Gln Ala Ala Ala Val Phe Ser Leu Val Asn			
1 5 10 15			
Glu Ile Phe Asn Arg Ser Ile Asn Leu Ile Val Ala Glu Leu Arg Leu			
20 25 30			
Gln Leu Asn Ala Arg Ala Glu Leu Asn Asn Leu Gln Arg Thr Leu Leu			
35 40 45			

Arg Thr His Ser Leu Leu Glu Glu Ala Lys Ala Arg Arg Met Thr Asp
50 55 60

Lys Ser Leu Val Leu Trp Leu Met Glu Leu Lys Glu Trp Ala Tyr Asp
65 70 75 80

Ala Asp Asp Ile Leu Asp Glu Tyr Glu Ala Ala Ala Ile Arg Leu Lys
85 90 95

Val Thr Arg Ser Thr Phe Lys Arg Leu Ile Asp His Val Ile Ile Asn
100 105 110

Val Pro Leu Ala His Lys Val Ala Asp Ile Arg Lys Arg Leu Asn Gly
115 120 125

Val Thr Leu Glu Arg Glu Leu Asn Leu Gly Ala Leu Glu Gly Ser Gln
130 135 140

Pro Leu Asp Ser Thr Lys Arg Gly Val Thr Thr Ser Leu Leu Thr Glu
145 150 155 160

Ser Cys Ile Val Gly Arg Ala Gln Asp Lys Glu Asn Leu Ile Arg Leu
165 170 175

Leu Leu Glu Pro Ser Asp Gly Ala Val Pro Val Val Pro Ile Val Gly
180 185 190

Leu Gly Gly Ala Gly Lys Thr Thr Leu Ser Gln Leu Ile Phe Asn Asp
195 200 205

Lys Arg Val Glu Glu His Phe Pro Leu Arg Met Trp Val Cys Val Ser
210 215 220

Asp Asp Phe Asp Val Lys Arg Ile Thr Arg Glu Ile Thr Glu Tyr Ala
225 230 235 240

Thr Asn Gly Arg Phe Met Asp Leu Thr Asn Leu Asn Met Leu Gln Val
245 250 255

Asn Leu Lys Glu Glu Ile Arg Gly Thr Thr Phe Leu Leu Val Leu Asp
260 265 270

Asp Val Trp Asn Glu Asp Pro Val Lys Trp Glu Ser Leu Leu Ala Pro
275 280 285

Leu Asp Ala Gly Gly Arg Gly Ser Val Val Ile Val Thr Thr Gln Ser
290 295 300

Lys Lys Val Ala Asp Val Thr Gly Thr Met Glu Pro Tyr Val Leu Glu
305 310 315 320
Glu Leu Thr Glu Asp Asp Ser Trp Ser Leu Ile Glu Ser His Ser Phe
325 330 335
Arg Glu Ala Ser Cys Ser Ser Thr Asn Pro Arg Met Glu Glu Ile Gly
340 345 350
Arg Lys Ile Ala Lys Lys Ile Ser Gly Leu Pro Tyr Gly Ala Thr Ala
355 360 365
Met Gly Arg Tyr Leu Arg Ser Lys His Gly Glu Ser Ser Trp Arg Glu
370 375 380
Val Leu Glu Thr Glu Thr Trp Glu Met Pro Pro Ala Ala Ser Asp Val
385 390 395 400
Leu Ser Ala Leu Arg Arg Ser Tyr Asp Asn Leu Pro Pro Gln Leu Lys
405 410 415
Leu Cys Phe Ala Phe Cys Ala Leu Phe Thr Lys Gly Tyr Arg Phe Arg
420 425 430
Lys Asp Thr Leu Ile His Met Trp Ile Ala Gln Asn Leu Ile Gln Ser
435 440 445
Thr Glu Ser Lys Arg Ser Glu Asp Met Ala Glu Glu Cys Phe Asp Asp
450 455 460
Leu Val Cys Arg Phe Phe Phe Arg Tyr Ser Trp Gly Asn Tyr Val Met
465 470 475 480
Asn Asp Ser Val His Asp Leu Ala Arg Trp Val Ser Leu Asp Glu Tyr
485 490 495
Phe Arg Ala Asp Glu Asp Ser Pro Leu His Ile Ser Lys Pro Ile Arg
500 505 510
His Leu Ser Trp Cys Ser Glu Arg Ile Thr Asn Val Leu Glu Asp Asn
515 520 525
Asn Thr Gly Gly Asp Ala Val Asn Pro Leu Ser Ser Leu Arg Thr Leu
530 535 540

Leu Phe Leu Gly Gln Ser Glu Phe Arg Ser Tyr His Leu Leu Asp Arg
 545 550 555 560

Met Phe Arg Met Leu Ser Arg Ile Arg Val Leu Asp Phe Ser Asn Cys
 565 570 575

Val Ile Arg Asn Leu Pro Ser Ser Val Gly Asn Leu Lys His Leu Arg
 580 585 590

Tyr Leu Gly Leu Ser Asn Thr Arg Ile Gln Arg Leu Pro Glu Ser Val
 595 600 605

Thr Arg Leu Cys Leu Leu Gln Thr Leu Leu Leu Glu Gly Cys Glu Leu
 610 615 620

Cys Arg Leu Pro Arg Ser Met Ser Arg Leu Val Lys Leu Arg Gln Leu
 625 630 635 640

Lys Ala Asn Pro Asp Val Ile Ala Asp Ile Ala Lys Val Gly Arg Leu
 645 650 655

Ile Glu Leu Gln Glu Leu Lys Ala Tyr Asn Val Asp Lys Lys Lys Gly
 660 665 670

His Gly Ile Ala Glu Leu Ser Ala Met Asn Gln Leu His Gly Asp Leu
 675 680 685

Ser Ile Arg Asn Leu Gln Asn Val Glu Lys Thr Arg Glu Ser Arg Lys
 690 695 700

Ala Arg Leu Asp Glu Lys Gln Lys Leu Lys Leu Leu Asp Leu Arg Trp
 705 710 715 720

Ala Asp Gly Arg Gly Ala Gly Glu Cys Asp Arg Asp Arg Lys Val Leu
 725 730 735

Lys Gly Leu Arg Pro His Pro Asn Leu Arg Glu Leu Ser Ile Lys Tyr
 740 745 750

Tyr Gly Gly Thr Ser Ser Pro Ser Trp Met Thr Asp Gln Tyr Leu Pro
 755 760 765

Asn Met Glu Thr Ile Arg Leu Arg Ser Cys Ala Arg Leu Thr Glu Leu
 770 775 780

Pro Cys Leu Gly Gln Leu His Ile Leu Arg His Leu His Ile Asp Gly
 785 790 795 800

Met Ser Gln Val Arg Gln Ile Asn Leu Gln Phe Tyr Gly Thr Gly Glu
 805 810 815

Val Ser Gly Phe Pro Leu Leu Glu Leu Leu Asn Ile Arg Arg Met Pro
 820 825 830

Ser Leu Glu Glu Trp Ser Glu Pro Arg Arg Asn Cys Cys Tyr Phe Pro
 835 840 845

Arg Leu His Lys Leu Leu Ile Glu Asp Cys Pro Arg Leu Arg Asn Leu
 850 855 860

Pro Ser Leu Pro Pro Thr Leu Glu Glu Leu Arg Ile Ser Arg Thr Gly
 865 870 875 880

Leu Val Asp Leu Pro Gly Phe His Gly Asn Gly Asp Val Thr Thr Asn
 885 890 895

Val Ser Leu Ser Ser Leu His Val Ser Glu Cys Arg Glu Leu Arg Ser
 900 905 910

Leu Ser Glu Gly Leu Leu Gln His Asn Leu Val Ala Leu Lys Thr Ala
 915 920 925

Ala Phe Thr Asp Cys Asp Ser Leu Glu Phe Leu Pro Ala Glu Gly Phe
 930 935 940

Arg Thr Ala Ile Ser Leu Glu Ser Leu Ile Met Thr Asn Cys Pro Leu
 945 950 955 960

Pro Cys Ser Phe Leu Leu Pro Ser Ser Leu Glu His Leu Lys Leu Gln
 965 970 975

Pro Cys Leu Tyr Pro Asn Asn Asn Glu Asp Ser Leu Ser Thr Cys Phe
 980 985 990

Glu Asn Leu Thr Ser Leu Ser Phe Leu Asp Ile Lys Asp Cys Pro Asn
 995 1000 1005

Leu Ser Ser Phe Pro Pro Gly Pro Leu Cys Gln Leu Ser Ala Leu
 1010 1015 1020

Gln His Leu Ser Leu Val Asn Cys Gln Arg Leu Gln Ser Ile Gly
 1025 1030 1035

Phe Gln Ala Leu Thr Ser Leu Glu Ser Leu Thr Ile Gln Asn Cys
1040 1045 1050

Pro Arg Leu Thr Met Ser His Ser Leu Val Glu Val Asn Asn Ser
1055 1060 1065

Ser Asp Thr Gly Leu Ala Phe Asn Ile Thr Arg Trp Met Arg Arg
1070 1075 1080

Arg Thr Gly Asp Asp Gly Leu Met Leu Arg His Arg Ala Gln Asn
1085 1090 1095

Asp Ser Phe Phe Gly Gly Leu Leu Gln His Leu Thr Phe Leu Gln
1100 1105 1110

Phe Leu Lys Ile Cys Gln Cys Pro Gln Leu Val Thr Phe Thr Gly
1115 1120 1125

Glu Glu Glu Glu Lys Trp Arg Asn Leu Thr Ser Leu Gln Ile Leu
1130 1135 1140

His Ile Val Asp Cys Pro Asn Leu Glu Val Leu Pro Ala Asn Leu
1145 1150 1155

Gln Ser Leu Cys Ser Leu Ser Thr Leu Tyr Ile Val Arg Cys Pro
1160 1165 1170

Arg Ile His Ala Phe Pro Pro Gly Gly Val Ser Met Ser Leu Ala
1175 1180 1185

His Leu Val Ile His Glu Cys Pro Gln Leu Cys Gln Arg Cys Asp
1190 1195 1200

Pro Pro Gly Gly Asp Asp Trp Pro Leu Ile Ala Asn Val Pro Arg
1205 1210 1215

Ile Cys Leu Gly Arg Thr His Pro Cys Arg Cys Ser Thr Thr
1220 1225 1230




FIGURE 1

		10	20	30	40	50	60	
RGA5	1	MSTALVIGGWFAQSFQTLDDKASNCATQCLARRGLHDDERRLRRTSLLRIEATLDKAEI						60
RGA2	1	-MADVTPQAAAVFSLVNEIFNRSINLIVAEELRLQLNARAEELNNLQRTLLRTHSLLEBAKA						59
		70	80	90	100	110	120	
RGA5	61	RWNHKNTSLVELVROLKDAAYDAEDLLESELYQAAKQKVEHRGQISDLFSESLSTASEW						120
RGA2	60	RR-MTDKSLVLWLMELKEWAYDADDILDEVEAAAIKLVTR---S---TEKRLIDHVI						110
		130	140	150	160	170	180	
RGA5	121	LGADGDDAGTRIREITQKILCNIAADMDVMQLLAPDDGGROFDWKVVERETSSFLTETVV						180
RGA2	111	INVP---LAHKVADIRKRLNGVTLEBELNLGALE---GSQPLDS-TKRGVITSLTESCT						163
		190	200	210	220	230	240	
RGA5	181	FGRDOEREKVVVELLLDSGSSNSSFSVLPLVGIGGVGKTTLAQLVYNDNRVGNVYFHLKVV						240
RGA2	164	VGRACDKENLIRLLLEPSDG--AVEVVPIVGLGAGKTTLSQLIFNDKRVEEHFFLRMAV						221
		250	260	270	280	290	300	
RGA5	241	CVSDNENVKRLTKEIESATKVEQSDKLNLDLQQLKEKTIASERFLVLDDVWSENRDD						300
RGA2	222	CVSDDFDVKRITREITEYATNGRFMDLTNLMLQVNLKEEIRGTTFLVLDDVWNEPDK						281
		310	320	330	340	350	360	
RGA5	301	WERLCAPLRFAARGSKVIVTTRDTKIASITCTMKEISLDGLQDDAYWELFKKCAFGSVN-						359
RGA2	282	WESILAPLDAGGRGSVVIVTTQSKKVADVTGTMEPYVLEELTEDDSWILIESHSFREASC						341
		370	380	390	400	410	420	
RGA5	360	FOEHLETLEVICRKIAGKIKGSLAAKTLCSLLRLDVSQEHRTIMESEVWOLPOAENEIL						419
RGA2	342	SSTNPRMEELGRKTAKKISCLPYGATAMGRYLRSKHGESSWREVLETETWEMPPAASDVL						401
		430	440	450	460	470	480	
RGA5	420	FVFWLSYQHLPGHILROCFAFCAVTHNDYLFYKHELTOIWLAEGETAHQGNKRMEDVGSSY						479
RGA2	402	SALRRSYDNLFPOLKLCFAPCALFTKCYRFRKDTLHMWIAQNLIQSTESKRSEDMAREC						461
		490	500	510	520	530	540	
RGA5	480	FHELVNRSFFQESRWGRGVVMDLIHDLAQFLSVGNCHRIDDDKSKETPSTTRHLSVALT						539
RGA2	462	EDDLVCRFFFRYS--WGVYVMNCSVHDLARWVSLDEYFRADEDSPLHISKPIRHLWCSE						519
		550	560	570	580	590	600	
RGA5	540	EQMCLVDFSGYN-KLRTLMINNQRNQPYMTKVNSCLPHSLFKRLKRIHVLVLOKCGMK						598
RGA2	520	RIITNVLEDNNTGGDAVNPISSLRTLLELGQSEFRSYHLLDRMERMLSRIRVLDHNSNCVIR						579

FIGURE 2-1

		610	620	630	640	650	660	
RGA5	599	ELPDILGDLIQLRYLDISYNAC	QRLPESL	CDLYNLOALRLWGCOLRSFPQMSKLINLR				658
RGA2	580	NLPSSVGNLKLRLRYGLS	NTRIQRLPESVTRCL	LOTLLEEGCEL	CLERSMSRLVKLR			638
		670	680	690	700	710	720	
RGA5	659	QLRVEDEILISKIYEVCKLISLOELSAFKVLNNHCNKLAEISGLTQIRSTLRITINLENVGS						718
RGA2	639	QLKANPDVIAITAKVGRLLIELOELKAYNVDKKKGCTAEISAMNQHGDLSIRNLQNVK						698
		730	740	750	760	770	780	
RGA5	719	KKEASKAKLHRKOYLEALELEWAAGQVSSLEHELIVSEEVLLGLOPHHFKSLITIRGYS						778
RGA2	699	TRESRKARLDEKOKIKLLDLRWADGRG	AGECDRDRKVLKGLRPHNIRELSIKYYGC					755
		790	800	810	820	830	840	
RGA5	779	ATVPSMIDVKMLPNLGLTLKLENC	TRLEGLSYICQLPHLRVLEMKRMPEVVKOMSHELCGCT					838
RGA2	756	TSSPSWMTDQYLPNMETIRLRSCARLE	TELECLGOLHILRLHLDGMSQVROINLOFYGTG					815
		850	860	870	880	890	900	
RGA5	839	KSKLEERLEELVIEDNPTLEKFEFNLAQLPCLKIIHMKNMFAVKHIGRELYGDIESNCFLS						898
RGA2	816	EVSGFELIUELLNIRRMPSLEWS						840
		910	920	930	940	950	960	
RGA5	899	LEELVLQDMLTLEELPNLQQLPHLRKVIHMKNMSALKLIGRELCDSEKIWFPRLEVLVLK						958
RGA2	840					RNCCYFPR		849
		970	980	990	1000	1010	1020	
RGA5	959	NMLALEELPSLDNFRVSRFFASSVEVGHLFSATRNKWFPRLEELEIKGMLTPEELHSLE						1018
RGA2	849							851
		1030	1040	1050	1060	1070	1080	
RGA5	1019	KLPCLKVFRIKGLPAVKKIGHGLFDSTCQRECFPRLEDVLSDMPAWEESWAEREELFS						1078
RGA2	851							851
		1090	1100	1110	1120	1130	1140	
RGA5	1079	CLCRLEKTEOCPEKLRCLLPPIHSLIKLELNQVGLTG	PGLCKGICGCSSTRTASLSLHTI					1138
RGA2	851							905
		1150	1160	1170	1180	1190	1200	
RGA5	1139	KCPNLRNLGECLLSNHLPHTNATIRIWECAELLWLPVKR	REFITLENLSIRNCPKLMST					1198
RGA2	906	ECRELRSLSEGLLOHNVALKTAATDCDSLEFLPAEGFRTAISLESLIMITNCP						963

FIGURE 2-2

		1210	1220	1230	1240	1250	1260	
RGA5	1199	QCEENDL	LLLEPLIKALELGDG	---NLGKSLEGCLENLSSLTOLATSNCPYMVSLPREVM				1255
RGA2	963	-----FLLSSLEHLKIQCLYFNNNEDSLSTCFENLSTSFLDIKOCENLSSFP	PGPL					1017
		1270	1280	1290	1300	1310	1320	
RGA5	1256	LHKKELGTVRDENCGLGSIEGLQVLKSLKRLAIIICCPRIILNR						1308
RGA2	1018	COLSALQHLISLVNCRLOSIG-FQALTSLESLLTONCPRLTMSHSLVEVNNS	SDTGLAFN					1076
		1330	1340	1350	1360	1370	1380	
RGA5	1309	LLLELSVDKT---	ALLKLSLIK	---TUPFIHSLRTIWSPOKVMFDLEEQLVHS				1356
RGA2	1077	ITRWMMRRRTGDDGLMLRHRAQND	SFFGGLLOHE	TELOFLKTCQCPOLVTF	TGEEEEKWRN			1136
		1390	1400	1410	1420	1430	1440	
RGA5	1357	LTALRRLEFRCKNLQSLPTELHTLPSTHALVWSDCPQIQSLEPEKGLPTL	ETDLGFDHCH					1416
RGA2	1137	LTSLOIILHTVDCPNIEVLPA	NLOSLSLSLYTVRCPR	IHAFFPGGVSM	SLAHVIHECP			1196
		1450	1460	1470				
RGA5	1417	PVLT-----	AQLEKHLAEMK--SSGRFHEVYA---					1441
RGA2	1197	QLCQRCDPPGGDDWPLIANVPRICLGRTHPCRCSTT						1232

FIGURE 2-3




FIGURE 3